

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
27 September 2001 (27.09.2001)

PCT

(10) International Publication Number  
**WO 01/70816 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/705**

MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(21) International Application Number: PCT/US01/09050

(22) International Filing Date: 21 March 2001 (21.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/191,355 22 March 2000 (22.03.2000) US  
60/269,799 20 February 2001 (20.02.2001) US

(71) Applicant (for all designated States except US): **ROHM AND HAAS COMPANY** [US/US]; 100 Independence Mall West, Philadelphia, PA 19106-2399 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PALLI, Subba, Reddy** [CA/US]; 832 Garfield Avenue, Lansdale, PA 19446 (US). **KAPITSKAYA, Marianna Zinovjevna** [RU/US]; 59 Wexford Road, North Wales, PA 19454 (US). **CRESS, Dean, Ervin** [US/US]; 104 Leslie Lane, Sounderton, PA 18964 (US).

(74) Agents: **MILLER, Suzanne, E.** et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations except US
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only
- of inventorship (Rule 4.17(iv)) for US only

#### Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL ECDYSONE RECEPTOR-BASED INDUCIBLE GENE EXPRESSION SYSTEM

**GAL4CfEcR**

**GAL4DNARD**

**CfEcRDEF**

**VP16RXR**

**VP16D**

**MmRXRDEF**

**pGAL4RELucGAL4RE TATA**

(57) Abstract: This invention relates to the field of biotechnology or genetic engineering. Specifically, this invention relates to the field of gene expression. More specifically, this invention relates to a novel inducible gene expression system and methods of modulating gene expression in a host cell for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic plants and animals.

## NOVEL ECDYSONE RECEPTOR-BASED INDUCIBLE GENE EXPRESSION SYSTEM

This application claims priority to co-pending US provisional application Serial  
5 number 60/191,355, filed March 22, 2000 and to co-pending US provisional application Serial  
number 60/269,799, filed February 20, 2001.

### FIELD OF THE INVENTION

10 This invention relates to the field of biotechnology or genetic engineering. Specifically,  
this invention relates to the field of gene expression. More specifically, this invention relates to  
a novel ecdysone receptor-based inducible gene expression system and methods of modulating  
the expression of a gene within a host cell using this inducible gene expression system.

### BACKGROUND OF THE INVENTION

In the field of genetic engineering, precise control of gene expression is a valuable tool  
for studying, manipulating, and controlling development and other physiological processes.  
Gene expression is a complex biological process involving a number of specific protein-protein  
20 interactions. In order for gene expression to be triggered, such that it produces the RNA  
necessary as the first step in protein synthesis, a transcriptional activator must be brought into  
proximity of a promoter that controls gene transcription. Typically, the transcriptional  
activator itself is associated with a protein that has at least one DNA binding domain that binds  
to DNA binding sites present in the promoter regions of genes. Thus, for gene expression to  
25 occur, a protein comprising a DNA binding domain and a transactivation domain located at an  
appropriate distance from the DNA binding domain must be brought into the correct position in  
the promoter region of the gene.

The traditional transgenic approach utilizes a cell-type specific promoter to drive the  
expression of a designed transgene. A DNA construct containing the transgene is first  
30 incorporated into a host genome. When triggered by a transcriptional activator, expression of  
the transgene occurs in a given cell type.

Another means to regulate expression of foreign genes in cells is through inducible  
promoters. Examples of the use of such inducible promoters include the PR1-a promoter,  
prokaryotic repressor-operator systems, immunosuppressive-immunophilin systems, and higher

eukaryotic transcription activation systems such as steroid hormone receptor systems and are described below.

The PR1-a promoter from tobacco is induced during the systemic acquired resistance response following pathogen attack. The use of PR1-a may be limited because it often responds to endogenous materials and external factors such as pathogens, UV-B radiation, and pollutants. Gene regulation systems based on promoters induced by heat shock, interferon and heavy metals have been described (Wurn et al., 1986, Proc. Natl. Acad. Sci. USA 83:5414-5418; Arnheiter et al., 1990 Cell 62:51-61; Filmus et al., 1992 Nucleic Acids Research 20:27550-27560). However, these systems have limitations due to their effect on expression of non-target genes. These systems are also leaky.

Prokaryotic repressor-operator systems utilize bacterial repressor proteins and the unique operator DNA sequences to which they bind. Both the tetracycline ("Tet") and lactose ("Lac") repressor-operator systems from the bacterium *Escherichia coli* have been used in plants and animals to control gene expression. In the Tet system, tetracycline binds to the TetR repressor protein, resulting in a conformational change which releases the repressor protein from the operator which as a result allows transcription to occur. In the Lac system, a lac operon is activated in response to the presence of lactose, or synthetic analogs such as isopropyl-b-D-thiogalactoside. Unfortunately, the use of such systems is restricted by unstable chemistry of the ligands, *i.e.* tetracycline and lactose, their toxicity, their natural presence, or the relatively high levels required for induction or repression. For similar reasons, utility of such systems in animals is limited.

Immunosuppressive molecules such as FK506, rapamycin and cyclosporine A can bind to immunophilins FKBP12, cyclophilin, *etc.* Using this information, a general strategy has been devised to bring together any two proteins simply by placing FK506 on each of the two proteins or by placing FK506 on one and cyclosporine A on another one. A synthetic homodimer of FK506 (FK1012) or a compound resulted from fusion of FK506-cyclosporine (FKCsA) can then be used to induce dimerization of these molecules (Spencer et al., 1993, *Science* 262:1019-24; Belshaw et al., 1996 *Proc Natl Acad Sci U S A* 93:4604-7). Gal4 DNA binding domain fused to FKBP12 and VP16 activator domain fused to cyclophilin, and FKCsA compound were used to show heterodimerization and activation of a reporter gene under the control of a promoter containing Gal4 binding sites. Unfortunately, this system includes immunosuppressants that can have unwanted side effects and therefore, limits its use for various mammalian gene switch applications.

Higher eukaryotic transcription activation systems such as steroid hormone receptor systems have also been employed. Steroid hormone receptors are members of the nuclear receptor superfamily and are found in vertebrate and invertebrate cells. Unfortunately, use of steroidal compounds that activate the receptors for the regulation of gene expression, particularly in plants and mammals, is limited due to their involvement in many other natural biological pathways in such organisms. In order to overcome such difficulties, an alternative system has been developed using insect ecdysone receptors (EcR).

Growth, molting, and development in insects are regulated by the ecdysone steroid hormone (molting hormone) and the juvenile hormones (Dhadialla, et al., 1998. Annu. Rev. Entomol. 43: 545-569). The molecular target for ecdysone in insects consists of at least ecdysone receptor (EcR) and ultraspiracle protein (USP). EcR is a member of the nuclear steroid receptor super family that is characterized by signature DNA and ligand binding domains, and an activation domain (Koelle et al. 1991, Cell, 67:59-77). EcR receptors are responsive to a number of steroidal compounds such as ponasterone A and muristerone A. Recently, non-steroidal compounds with ecdysteroid agonist activity have been described, including the commercially available insecticides tebufenozide and methoxyfenozide that are marketed world wide by Rohm and Haas Company (see International Patent Application No. PCT/EP96/00686 and US Patent 5,530,028). Both analogs have exceptional safety profiles to other organisms.

International Patent Application No. PCT/US97/05330 (WO 97/38117) discloses methods for modulating the expression of an exogenous gene in which a DNA construct comprising the exogenous gene and an ecdysone response element is activated by a second DNA construct comprising an ecdysone receptor that, in the presence of a ligand therefor, and optionally in the presence of a receptor capable of acting as a silent partner, binds to the ecdysone response element to induce gene expression. The ecdysone receptor of choice was isolated from *Drosophila melanogaster*. Typically, such systems require the presence of the silent partner, preferably retinoid X receptor (RXR), in order to provide optimum activation. In mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and regulates expression of target genes in a ligand dependent manner. International Patent Application No. PCT/US98/14215 (WO 99/02683) discloses that the ecdysone receptor isolated from the silk moth *Bombyx mori* is functional in mammalian systems without the need for an exogenous dimer partner.

U.S. Patent No. 5,880,333 discloses a *Drosophila melanogaster* EcR and ultraspiracle



(USP) heterodimer system used in plants in which the transactivation domain and the DNA binding domain are positioned on two different hybrid proteins. Unfortunately, this system is not effective for inducing reporter gene expression in animal cells (for comparison, see Example 1.2, below).

5 In each of these cases, the transactivation domain and the DNA binding domain (either as native EcR as in International Patent Application No. PCT/US98/14215 or as modified EcR as in International Patent Application No. PCT/US97/05330) were incorporated into a single molecule and the other heterodimeric partners, either USP or RXR, were used in their native state.

10 Drawbacks of the above described EcR-based gene regulation systems include a considerable background activity in the absence of ligands and that these systems are not applicable for use in both plants and animals (see U.S. Patent No. 5,880,333). For most applications that rely on modulating gene expression, these EcR-based systems are undesirable. Therefore, a need exists in the art for improved systems to precisely modulate the expression of  
15 exogenous genes in both plants and animals. Such improved systems would be useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic animals. Improved systems that are simple, compact, and dependent on ligands that are relatively inexpensive, readily available, and of low toxicity to the host would prove useful for  
20 regulating biological systems.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties. However, the citation of any reference herein should not be construed as an admission that such reference is available as "Prior Art" to the instant application.

25

## **SUMMARY OF THE INVENTION**

The present invention relates to a novel ecdysone receptor-based inducible gene expression system, novel receptor polynucleotides and polypeptides for use in the novel  
30 inducible gene expression system, and methods of modulating the expression of a gene within a host cell using this inducible gene expression system. In particular, Applicants' invention relates to an improved gene expression modulation system comprising a polynucleotide encoding a receptor polypeptide comprising a truncation mutation.

Specifically, the present invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide that encodes a first polypeptide comprising: i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than an ultraspiracle receptor; wherein the DNA binding domain and the transactivation domain are from a polypeptide other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor; wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor (EcR) ligand binding domain

In another specific embodiment, the ligand binding domain of the second polypeptide comprises a retinoid X receptor (RXR) ligand binding domain.

In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain

The present invention also relates to a gene expression modulation system according to the invention further comprising c) a third gene expression cassette comprising: i) a response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

The present invention also relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide, wherein the truncation mutation affects ligand binding activity or ligand sensitivity.

In particular, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific

embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

The present invention also relates to an isolated polynucleotide encoding a truncated  
5 EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of said EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of said EcR or RXR polypeptide. In another specific  
10 embodiment, the present invention relates to an isolated polynucleotide encoding a truncated EcR or a truncated RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of said EcR or RXR polypeptide.

The present invention also relates to an isolated polynucleotide encoding a truncated RXR polypeptide comprising a truncation mutation that increases ligand sensitivity of a  
15 heterodimer comprising the truncated retinoid X receptor polypeptide and a dimerization partner. In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide.

The present invention also relates to an isolated polypeptide encoded by a polynucleotide according to Applicants' invention. In particular, the present invention relates  
20 to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation, wherein the EcR or RXR polypeptide is encoded by a polynucleotide according to the invention.

Thus, the present invention also relates to an isolated truncated EcR or truncated RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand  
25 sensitivity of said EcR or RXR polypeptide.

Applicants' invention also relates to methods of modulating gene expression in a host cell using a gene expression modulation system according to the invention. Specifically, Applicants' invention provides a method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of: a) introducing into the host cell a  
30 gene expression modulation system according to the invention; and b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide of the gene expression modulation system; wherein the gene to be expressed is a component of a chimeric gene comprising: i) a response element

comprising a domain to which the DNA binding domain from the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated, whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates

5 expression of the gene in the host cell.

Applicants' invention also provides an isolated host cell comprising an inducible gene expression system according to the invention. The present invention also relates to an isolated host cell comprising a polynucleotide or polypeptide according to the invention. Accordingly, Applicants' invention also relates to a non-human organism comprising a host cell according to

10 the invention.

**BRIEF DESCRIPTION OF THE DRAWINGS**

**Figure 1:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-MmRXRDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.1).

**Figure 2:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfUSPDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.2).

**Figure 3:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.3).

**Figure 4:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-MmRXRDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-DmEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.4).

**Figure 5:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfUSPDEF chimeric polypeptide and a second gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.5).

**Figure 6:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a Gal4DBD-CfEcRDEF-VP16AD chimeric polypeptide; prepared as described in Example 1 (switch 1.6).

**Figure 7:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.7).

**Figure 8:** An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-DmEcRCDEF chimeric polypeptide and a second gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.8).

Figure 9: An ecdysone receptor-based gene expression system comprising a first gene expression cassette encoding a VP16AD-CfEcRCDEF chimeric polypeptide and a second gene expression cassette encoding a MmRXR polypeptide; prepared as described in Example 1 (switch 1.9).

- 5 Figure 10: An ecdysone receptor-based gene expression system comprising a gene expression cassette encoding a Gal4DBD-CfEcRCDEF chimeric polypeptide; prepared as described in Example 1 (switch 1.10).

Figure 11: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation  
10 mutants transfected into NIH3T3 cells along with VP16MmRXRDE, pFRLUc and pTKRL plasmid DNAs.

Figure 12: Expression data of GAL4CfEcRA/BCDEF, GAL4CfEcRCDEF, GAL4CfEcR1/2CDEF, GAL4CfEcRDEF, GAL4CfEcREF, GAL4CfEcRDE truncation  
15 mutants transfected into 3T3 cells along with VP16MmRXRE, pFRLUc and pTKRL plasmid DNAs.

Figure 13: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del  
constructs transfected into NIH3T3 cells along with GAL4CfEcRCDEF, pFRLUc and pTKRL  
plasmid DNAs.

20 Figure 14: Expression data of VP16MmRXRA/BCDEF, VP16MmRXRCDEF, VP16MmRXRDEF, VP16MmRXREF, VP16MmRXRBam-EF, VP16MmRXRAF2del  
constructs transfected into NIH3T3 cells along with GAL4CfEcRDEF, pFRLUc and pTKRL  
plasmid DNAs.

Figure 15: Expression data of various truncated CfEcR and MmRXR receptor pairs  
25 transfected into NIH3T3 cells along with GAL4CfEcRDEF, pFRLUc and pTKRL plasmid  
DNAs.

### **DETAILED DESCRIPTION OF THE INVENTION**

30 Applicants have now developed an improved ecdysone receptor-based, inducible gene expression system comprising a truncation mutant of an ecdysone receptor or a retinoid X receptor (RXR) polypeptide that affects ligand binding activity or ligand sensitivity. This mutational effect may increase or reduce ligand binding activity or ligand sensitivity and may

be steroid or non-steroid specific. Thus, Applicants' invention provides an improved ecdysone receptor-based inducible gene expression system useful for modulating expression of a gene of interest in a host cell. In a particularly desirable embodiment, Applicants' invention provides an inducible gene expression system that has a reduced level of background gene expression and responds to submicromolar concentrations of non-steroidal ligand. Thus, Applicants' novel inducible gene expression system and its use in methods of modulating gene expression in a host cell overcome the limitations of currently available inducible expression systems and provide the skilled artisan with an effective means to control gene expression.

The present invention provides a novel inducible gene expression system that can be used to modulate gene expression in both prokaryotic and eukaryotic host cells. Applicants' invention is useful for applications such as gene therapy, large scale production of proteins and antibodies, cell-based high throughput screening assays, functional genomics and regulation of traits in transgenic organisms.

## 15 DEFINITIONS

In this disclosure, a number of terms and abbreviations are used. The following definitions are provided and should be helpful in understanding the scope and practice of the present invention.

In a specific embodiment, the term "about" or "approximately" means within 20%, preferably within 10%, more preferably within 5%, and even more preferably within 1% of a given value or range.

The term "substantially free" means that a composition comprising "A" (where "A" is a single protein, DNA molecule, vector, recombinant host cell, etc.) is substantially free of "B" (where "B" comprises one or more contaminating proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight. It is also preferred that a composition, which is substantially free of contamination, contain only a single molecular weight species having the activity or characteristic of the species of interest.

The term "isolated" for the purposes of the present invention designates a biological material (nucleic acid or protein) that has been removed from its original environment (the environment in which it is naturally present).

For example, a polynucleotide present in the natural state in a plant or an animal is not isolated. The same polynucleotide separated from the adjacent nucleic acids in which it is naturally present. The term "purified" does not require the material to be present in a form exhibiting absolute purity, exclusive of the presence of other compounds. It is rather a relative  
5 definition.

A polynucleotide is in the "purified" state after purification of the starting material or of the natural material by at least one order of magnitude, preferably 2 or 3 and preferably 4 or 5 orders of magnitude.

A "nucleic acid" is a polymeric compound comprised of covalently linked subunits  
10 called nucleotides. Nucleic acid includes polyribonucleic acid (RNA) and polydeoxyribonucleic acid (DNA), both of which may be single-stranded or double-stranded. DNA includes but is not limited to cDNA, genomic DNA, plasmids DNA, synthetic DNA, and semi-synthetic DNA. DNA may be linear, circular, or supercoiled.

A "nucleic acid molecule" refers to the phosphate ester polymeric form of  
15 ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in  
20 particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal  
25 convention of giving only the sequence in the 5' to 3' direction along the non-transcribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

The term "fragment" will be understood to mean a nucleotide sequence of reduced length relative to the reference nucleic acid and comprising, over the common portion, a  
30 nucleotide sequence identical to the reference nucleic acid. Such a nucleic acid fragment according to the invention may be, where appropriate, included in a larger polynucleotide of which it is a constituent. Such fragments comprise, or alternatively consist of, oligonucleotides ranging in length from at least 8, 10, 12, 15, 18, 20 to 25, 30, 40, 50, 70, 80, 100, 200, 500,



1000 or 1500 consecutive nucleotides of a nucleic acid according to the invention.

As used herein, an "isolated nucleic acid fragment" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised  
5 of one or more segments of cDNA, genomic DNA or synthetic DNA.

A "gene" refers to an assembly of nucleotides that encode a polypeptide, and includes cDNA and genomic DNA nucleic acids. "Gene" also refers to a nucleic acid fragment that expresses a specific protein or polypeptide, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene"  
10 refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and/or coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that  
15 found in nature. A chimeric gene may comprise coding sequences derived from different sources and/or regulatory sequences derived from different sources. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene or "heterologous" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes  
20 inserted into a non-native organism, or chimeric genes. A "transgene" is a gene that has been introduced into the genome by a transformation procedure.

"Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

25 The term "genome" includes chromosomal as well as mitochondrial, chloroplast and viral DNA or RNA.

A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and  
30 solution ionic strength (see Sambrook *et al.*, 1989 *infra*). Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly Chapter 11 and Table 11.1 therein (entirely

incorporated herein by reference). The conditions of temperature and ionic strength determine the "stringency" of the hybridization.

Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as  
5 genes that duplicate functional enzymes from closely related organisms. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a  $T_m$  of  $55^\circ$ , can be used, *e.g.*, 5x SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5x SSC, 0.5% SDS). Moderate stringency hybridization conditions correspond to a higher  $T_m$ , *e.g.*, 40% formamide, with 5x or 6x SCC. High stringency hybridization  
10 conditions correspond to the highest  $T_m$ , *e.g.*, 50% formamide, 5x or 6x SCC. Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

The term "complementary" is used to describe the relationship between nucleotide bases that are capable of hybridizing to one another. For example, with respect to DNA,  
15 adenosine is complementary to thymine and cytosine is complementary to guanine. Accordingly, the instant invention also includes isolated nucleic acid fragments that are complementary to the complete sequences as disclosed or used herein as well as those substantially similar nucleic acid sequences.

In a specific embodiment, the term "standard hybridization conditions" refers to a  $T_m$   
20 of  $55^\circ\text{C}$ , and utilizes conditions as set forth above. In a preferred embodiment, the  $T_m$  is  $60^\circ\text{C}$ ; in a more preferred embodiment, the  $T_m$  is  $65^\circ\text{C}$ .

Post-hybridization washes also determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 minutes (min), then repeated with 2X SSC, 0.5% SDS at  $45^\circ\text{C}$  for 30 minutes, and then  
25 repeated twice with 0.2X SSC, 0.5% SDS at  $50^\circ\text{C}$  for 30 minutes. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to  $60^\circ\text{C}$ . Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at  $65^\circ\text{C}$ . Hybridization requires that the two nucleic acids comprise  
30 complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible.

The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater

the degree of similarity or homology between two nucleotide sequences, the greater the value of  $T_m$  for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher  $T_m$ ) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating  $T_m$  have been derived (see Sambrook *et al.*, *supra*, 9.50-0.51). For hybridization with shorter nucleic acids, *i.e.*, oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook *et al.*, *supra*, 11.7-11.8).

In one embodiment the length for a hybridizable nucleic acid is at least about 10 nucleotides. Preferable a minimum length for a hybridizable nucleic acid is at least about 15 nucleotides; more preferably at least about 20 nucleotides; and most preferably the length is at least 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

The term "probe" refers to a single-stranded nucleic acid molecule that can base pair with a complementary single stranded target nucleic acid to form a double-stranded molecule.

As used herein, the term "oligonucleotide" refers to a nucleic acid, generally of at least 18 nucleotides, that is hybridizable to a genomic DNA molecule, a cDNA molecule, a plasmid DNA or an mRNA molecule. Oligonucleotides can be labeled, *e.g.*, with  $^{32}\text{P}$ -nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. A labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid.

Oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of a nucleic acid, or to detect the presence of a nucleic acid.

An oligonucleotide can also be used to form a triple helix with a DNA molecule. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer.

Accordingly, oligonucleotides can be prepared with non-naturally occurring phosphoester analog bonds, such as thioester bonds, etc.

A "primer" is an oligonucleotide that hybridizes to a target nucleic acid sequence to create a double stranded nucleic acid region that can serve as an initiation point for DNA synthesis under suitable conditions. Such primers may be used in a polymerase chain reaction.

"Polymerase chain reaction" is abbreviated PCR and means an *in vitro* method for enzymatically amplifying specific nucleic acid sequences. PCR involves a repetitive series of temperature cycles with each cycle comprising three stages: denaturation of the template

nucleic acid to separate the strands of the target molecule, annealing a single stranded PCR oligonucleotide primer to the template nucleic acid, and extension of the annealed primer(s) by DNA polymerase. PCR provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

“Reverse transcription-polymerase chain reaction” is abbreviated RT-PCR and means an *in vitro* method for enzymatically producing a target cDNA molecule or molecules from an RNA molecule or molecules, followed by enzymatic amplification of a specific nucleic acid sequence or sequences within the target cDNA molecule or molecules as described above. RT-PCR also provides a means to detect the presence of the target molecule and, under quantitative or semi-quantitative conditions, to determine the relative amount of that target molecule within the starting pool of nucleic acids.

A DNA “coding sequence” is a double-stranded DNA sequence that is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. “Suitable regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, polyadenylation recognition sequences, RNA processing site, effector binding site and stem-loop structure. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from mRNA, genomic DNA sequences, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

“Open reading frame” is abbreviated ORF and means a length of nucleic acid sequence, either DNA, cDNA or RNA, that comprises a translation start signal or initiation codon, such as an ATG or AUG, and a termination codon and can be potentially translated into a polypeptide sequence.

The term “head-to-head” is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-head orientation when the 5' end of the coding strand of one polynucleotide is adjacent

to the 5' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds away from the 5' end of the other polynucleotide.

The term "head-to-head" may be abbreviated (5')-to-(5') and may also be indicated by the symbols ( $\leftarrow \rightarrow$ ) or ( $3' \leftarrow 5' 5' \rightarrow 3'$ ).

5        The term "tail-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a tail-to-tail orientation when the 3' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each polynucleotide proceeds toward the other polynucleotide. The term "tail-to-tail" may be  
10 abbreviated (3')-to-(3') and may also be indicated by the symbols ( $\rightarrow \leftarrow$ ) or ( $5' \rightarrow 3' 3' \leftarrow 5'$ ).

The term "head-to-tail" is used herein to describe the orientation of two polynucleotide sequences in relation to each other. Two polynucleotides are positioned in a head-to-tail orientation when the 5' end of the coding strand of one polynucleotide is adjacent to the 3' end of the coding strand of the other polynucleotide, whereby the direction of transcription of each  
15 polynucleotide proceeds in the same direction as that of the other polynucleotide. The term "head-to-tail" may be abbreviated (5')-to-(3') and may also be indicated by the symbols ( $\rightarrow \rightarrow$ ) or ( $5' \rightarrow 3' 5' \rightarrow 3'$ ).

The term "downstream" refers to a nucleotide sequence that is located 3' to reference nucleotide sequence. In particular, downstream nucleotide sequences generally relate to  
20 sequences that follow the starting point of transcription. For example, the translation initiation codon of a gene is located downstream of the start site of transcription.

The term "upstream" refers to a nucleotide sequence that is located 5' to reference nucleotide sequence. In particular, upstream nucleotide sequences generally relate to sequences that are located on the 5' side of a coding sequence or starting point of transcription. For  
25 example, most promoters are located upstream of the start site of transcription.

The terms "restriction endonuclease" and "restriction enzyme" refer to an enzyme that binds and cuts within a specific nucleotide sequence within double stranded DNA.

"Homologous recombination" refers to the insertion of a foreign DNA sequence into another DNA molecule, e.g., insertion of a vector in a chromosome. Preferably, the vector  
30 targets a specific chromosomal site for homologous recombination. For specific homologous recombination, the vector will contain sufficiently long regions of homology to sequences of the chromosome to allow complementary binding and incorporation of the vector into the chromosome. Longer regions of homology, and greater degrees of sequence similarity, may

increase the efficiency of homologous recombination.

Several methods known in the art may be used to propagate a polynucleotide according to the invention. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As described  
5 herein, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (*e.g.*, lambda), and plasmid and cosmid DNA vectors, to name but a few.

A "vector" is any means for the cloning of and/or transfer of a nucleic acid into a host  
10 cell. A vector may be a replicon to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (*e.g.*, plasmid, phage, cosmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, *i.e.*, capable of replication under its own control. The term "vector" includes both viral and nonviral means for introducing the nucleic acid into a cell *in vitro*, *ex vivo* or *in vivo*. A large number of vectors known in the art may be used to manipulate nucleic  
15 acids, incorporate response elements and promoters into genes, etc. Possible vectors include, for example, plasmids or modified viruses including, for example bacteriophages such as lambda derivatives, or plasmids such as PBR322 or pUC plasmid derivatives, or the Bluescript vector. For example, the insertion of the DNA fragments corresponding to response elements  
20 and promoters into a suitable vector can be accomplished by ligating the appropriate DNA fragments into a chosen vector that has complementary cohesive termini. Alternatively, the ends of the DNA molecules may be enzymatically modified or any site may be produced by ligating nucleotide sequences (linkers) into the DNA termini. Such vectors may be engineered to contain selectable marker genes that provide for the selection of cells that have incorporated  
25 the marker into the cellular genome. Such markers allow identification and/or selection of host cells that incorporate and express the proteins encoded by the marker.

Viral vectors, and particularly retroviral vectors, have been used in a wide variety of gene delivery applications in cells, as well as living animal subjects. Viral vectors that can be used include but are not limited to retrovirus, adeno-associated virus, pox, baculovirus,  
30 vaccinia, herpes simplex, Epstein-Barr, adenovirus, geminivirus, and caulimovirus vectors. Non-viral vectors include plasmids, liposomes, electrically charged lipids (cytofectins), DNA-protein complexes, and biopolymers. In addition to a nucleic acid, a vector may also comprise one or more regulatory regions, and/or selectable markers useful in selecting, measuring, and

monitoring nucleic acid transfer results (transfer to which tissues, duration of expression, etc.).

The term "plasmid" refers to an extra chromosomal element often carrying a gene that is not part of the central metabolism of the cell, and usually in the form of circular double-stranded DNA molecules. Such elements may be autonomously replicating sequences, genome  
5 integrating sequences, phage or nucleotide sequences, linear, circular, or supercoiled, of a single- or double-stranded DNA or RNA, derived from any source, in which a number of nucleotide sequences have been joined or recombined into a unique construction which is capable of introducing a promoter fragment and DNA sequence for a selected gene product along with appropriate 3' untranslated sequence into a cell.

10 A "cloning vector" is a "replicon", which is a unit length of a nucleic acid, preferably DNA, that replicates sequentially and which comprises an origin of replication, such as a plasmid, phage or cosmid, to which another nucleic acid segment may be attached so as to bring about the replication of the attached segment. Cloning vectors may be capable of replication in one cell type and expression in another ("shuttle vector").

15 Vectors may be introduced into the desired host cells by methods known in the art, *e.g.*, transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, *e.g.*, Wu et al., 1992, J. Biol. Chem. 267:963-967; Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624; and Hartmut et al., Canadian Patent Application No.  
20 2,012,311, filed March 15, 1990).

A polynucleotide according to the invention can also be introduced *in vivo* by lipofection. For the past decade, there has been increasing use of liposomes for encapsulation and transfection of nucleic acids *in vitro*. Synthetic cationic lipids designed to limit the difficulties and dangers encountered with liposome mediated transfection can be used to prepare liposomes for *in vivo*  
25 transfection of a gene encoding a marker (Felgner et al., 1987. PNAS 84:7413; Mackey, et al., 1988. Proc. Natl. Acad. Sci. U.S.A. 85:8027-8031; and Ulmer et al., 1993. Science 259:1745-1748). The use of cationic lipids may promote encapsulation of negatively charged nucleic acids, and also promote fusion with negatively charged cell membranes (Felgner and Ringold, 1989. Science 337:387-388). Particularly useful lipid compounds and compositions for transfer of  
30 nucleic acids are described in International Patent Publications WO95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127. The use of lipofection to introduce exogenous genes into the specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. It is clear that directing transfection to particular cell

types would be particularly preferred in a tissue with cellular heterogeneity, such as pancreas, liver, kidney, and the brain. Lipids may be chemically coupled to other molecules for the purpose of targeting (Mackey, et al., 1988, *supra*). Targeted peptides, *e.g.*, hormones or neurotransmitters, and proteins such as antibodies, or non-peptide molecules could be coupled to liposomes

5 chemically.

Other molecules are also useful for facilitating transfection of a nucleic acid *in vivo*, such as a cationic oligopeptide (*e.g.*, WO95/21931), peptides derived from DNA binding proteins (*e.g.*, WO96/25508), or a cationic polymer (*e.g.*, WO95/21931).

It is also possible to introduce a vector *in vivo* as a naked DNA plasmid (see U.S.  
10 Patents 5,693,622, 5,589,466 and 5,580,859). Receptor-mediated DNA delivery approaches can also be used (Curiel et al., 1992. Hum. Gene Ther. 3:147-154; and Wu and Wu, 1987. J. Biol. Chem. 262:4429-4432).

The term "transfection" means the uptake of exogenous or heterologous RNA or DNA by a cell. A cell has been "transfected" by exogenous or heterologous RNA or DNA when  
15 such RNA or DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous RNA or DNA when the transfected RNA or DNA effects a phenotypic change. The transforming RNA or DNA can be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.

"Transformation" refers to the transfer of a nucleic acid fragment into the genome of a  
20 host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" or "recombinant" or "transformed" organisms.

The term "genetic region" will refer to a region of a nucleic acid molecule or a nucleotide sequence that comprises a gene encoding a polypeptide.

25 In addition, the recombinant vector comprising a polynucleotide according to the invention may include one or more origins for replication in the cellular hosts in which their amplification or their expression is sought, markers or selectable markers.

The term "selectable marker" means an identifying factor, usually an antibiotic or chemical resistance gene, that is able to be selected for based upon the marker gene's effect,  
30 *i.e.*, resistance to an antibiotic, resistance to a herbicide, colorimetric markers, enzymes, fluorescent markers, and the like, wherein the effect is used to track the inheritance of a nucleic acid of interest and/or to identify a cell or organism that has inherited the nucleic acid of



interest. Examples of selectable marker genes known and used in the art include: genes providing resistance to ampicillin, streptomycin, gentamycin, kanamycin, hygromycin, bialaphos herbicide, sulfonamide, and the like; and genes that are used as phenotypic markers, *i.e.*, anthocyanin regulatory genes, isopentenyl transferase gene, and the like.

5           The term "reporter gene" means a nucleic acid encoding an identifying factor that is able to be identified based upon the reporter gene's effect, wherein the effect is used to track the inheritance of a nucleic acid of interest, to identify a cell or organism that has inherited the nucleic acid of interest, and/or to measure gene expression induction or transcription. Examples of reporter genes known and used in the art include: luciferase (Luc), green fluorescent protein  
10 (GFP), chloramphenicol acetyltransferase (CAT),  $\beta$ -galactosidase (LacZ),  $\beta$ -glucuronidase (Gus), and the like. Selectable marker genes may also be considered reporter genes.

          "Promoter" refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. Promoters may be derived in their entirety from a native gene, or be composed of  
15 different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental or physiological conditions. Promoters that cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive  
20 promoters". Promoters that cause a gene to be expressed in a specific cell type are commonly referred to as "cell-specific promoters" or "tissue-specific promoters". Promoters that cause a gene to be expressed at a specific stage of development or cell differentiation are commonly referred to as "developmentally-specific promoters" or "cell differentiation-specific promoters". Promoters that are induced and cause a gene to be expressed following exposure or treatment  
25 of the cell with an agent, biological molecule, chemical, ligand, light, or the like that induces the promoter are commonly referred to as "inducible promoters" or "regulatable promoters". It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of different lengths may have identical promoter activity.

30           A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include

the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

5 A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced (if the coding sequence contains introns) and translated into the protein encoded by the coding sequence.

“Transcriptional and translational control sequences” are DNA regulatory sequences,  
10 such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

The term “response element” means one or more cis-acting DNA elements which confer responsiveness on a promoter mediated through interaction with the DNA-binding  
15 domains of the first chimeric gene. This DNA element may be either palindromic (perfect or imperfect) in its sequence or composed of sequence motifs or half sites separated by a variable number of nucleotides. The half sites can be similar or identical and arranged as either direct or inverted repeats or as a single half site or multimers of adjacent half sites in tandem. The response element may comprise a minimal promoter isolated from different organisms  
20 depending upon the nature of the cell or organism into which the response element will be incorporated. The DNA binding domain of the first hybrid protein binds, in the presence or absence of a ligand, to the DNA sequence of a response element to initiate or suppress transcription of downstream gene(s) under the regulation of this response element. Examples of DNA sequences for response elements of the natural ecdysone receptor include:

25 RRGG/TTCANTGAC/ACY (see Cherbas L., et. al., (1991), *Genes Dev.* 5, 120-131);  
AGGTCAN<sub>(n)</sub>AGGTCA, where N<sub>(n)</sub> can be one or more spacer nucleotides (see D'Avino PP., et. al., (1995), *Mol. Cell. Endocrinol.* 113, 1-9); and GGGTTGAATGAATTT (see Antoniewski C., et. al., (1994). *Mol. Cell Biol.* 14, 4465-4474).

The term “operably linked” refers to the association of nucleic acid sequences on a  
30 single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in

sense or antisense orientation.

The term “expression”, as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from a nucleic acid or polynucleotide. Expression may also refer to translation of mRNA into a protein or  
5 polypeptide.

The terms “cassette”, “expression cassette” and “gene expression cassette” refer to a segment of DNA that can be inserted into a nucleic acid or polynucleotide at specific restriction sites or by homologous recombination. The segment of DNA comprises a polynucleotide that encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure  
10 insertion of the cassette in the proper reading frame for transcription and translation.

“Transformation cassette” refers to a specific vector comprising a polynucleotide that encodes a polypeptide of interest and having elements in addition to the polynucleotide that facilitate transformation of a particular host cell. Cassettes, expression cassettes, gene expression cassettes and transformation cassettes of the invention may also comprise elements that allow  
15 for enhanced expression of a polynucleotide encoding a polypeptide of interest in a host cell. These elements may include, but are not limited to: a promoter, a minimal promoter, an enhancer, a response element, a terminator sequence, a polyadenylation sequence, and the like.

For purposes of this invention, the term “gene switch” refers to the combination of a response element associated with a promoter, and an EcR based system which, in the presence  
20 of one or more ligands, modulates the expression of a gene into which the response element and promoter are incorporated.

The terms “modulate” and “modulates” mean to induce, reduce or inhibit nucleic acid or gene expression, resulting in the respective induction, reduction or inhibition of protein or polypeptide production.

25 The plasmids or vectors according to the invention may further comprise at least one promoter suitable for driving expression of a gene in a host cell. The term “expression vector” means a vector, plasmid or vehicle designed to enable the expression of an inserted nucleic acid sequence following transformation into the host. The cloned gene, i.e., the inserted nucleic acid sequence, is usually placed under the control of control elements such as a promoter, a minimal  
30 promoter, an enhancer, or the like. Initiation control regions or promoters, which are useful to drive expression of a nucleic acid in the desired host cell are numerous and familiar to those skilled in the art. Virtually any promoter capable of driving these genes is suitable for the present invention including but not limited to: viral promoters, plant promoters, bacterial

promoters, animal promoters, mammalian promoters, synthetic promoters, constitutive  
 promoters, tissue specific promoter, developmental specific promoters, inducible promoters,  
 light regulated promoters; *CYC1*, *HIS3*, *GAL1*, *GAL4*, *GAL10*, *ADH1*, *PGK*, *PHO5*, *GAPDH*,  
 5 *ADC1*, *TRP1*, *URA3*, *LEU2*, *ENO*, *TPI*, alkaline phosphatase promoters (useful for expression  
 in *Saccharomyces*); *AOX1* promoter (useful for expression in *Pichia*); b-lactamase, *lac*, *ara*,  
*tet*, *trp*, *lP<sub>L</sub>*, *lP<sub>R</sub>*, *T7*, *tac*, and *trc* promoters (useful for expression in *Escherichia coli*); and  
 light regulated-, seed specific-, pollen specific-, ovary specific-, pathogenesis or disease  
 related-, cauliflower mosaic virus 35S, CMV 35S minimal, cassava vein mosaic virus  
 (CsVMV), chlorophyll a/b binding protein, ribulose 1, 5-bisphosphate carboxylase, shoot-  
 10 specific, root specific, chitinase, stress inducible, rice tungro bacilliform virus, plant super-  
 promoter, potato leucine aminopeptidase, nitrate reductase, mannopine synthase, nopaline  
 synthase, ubiquitin, zein protein, and anthocyanin promoters (useful for expression in plant  
 cells); animal and mammalian promoters known in the art include, but are not limited to, the  
 SV40 early (SV40e) promoter region, the promoter contained in the 3' long terminal repeat  
 15 (LTR) of Rous sarcoma virus (RSV), the promoters of the E1A or major late promoter (MLP)  
 genes of adenoviruses, the cytomegalovirus early promoter, the herpes simplex virus (HSV)  
 thymidine kinase (TK) promoter, an elongation factor 1 alpha (EF1) promoter, a  
 phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, an albumin promoter,  
 the regulatory sequences of the mouse metallothionein-L promoter, and transcriptional control  
 20 regions, the ubiquitous promoters (HPRT, vimentin,  $\alpha$ -actin, tubulin and the like), the  
 promoters of the intermediate filaments (desmin, neurofilaments, keratin, GFAP, and the like),  
 the promoters of therapeutic genes (of the MDR, CFTR or factor VIII type, and the like), and  
 promoters that exhibit tissue specificity and have been utilized in transgenic animals, such as  
 the elastase I gene control region which is active in pancreatic acinar cells; insulin gene control  
 25 region active in pancreatic beta cells, immunoglobulin gene control region active in lymphoid  
 cells, mouse mammary tumor virus control region active in testicular, breast, lymphoid and  
 mast cells; albumin gene, Apo AI and Apo AII control regions active in liver, alpha-fetoprotein  
 gene control region active in liver, alpha 1-antitrypsin gene control region active in the liver,  
 beta-globin gene control region active in myeloid cells, myelin basic protein gene control region  
 30 active in oligodendrocyte cells in the brain, myosin light chain-2 gene control region active in  
 skeletal muscle, and gonadotropic releasing hormone gene control region active in the  
 hypothalamus, pyruvate kinase promoter, villin promoter, promoter of the fatty acid binding  
 intestinal protein, promoter of the smooth muscle cell  $\alpha$ -actin, and the like. In a preferred

embodiment of the invention, the promoter is selected from the group consisting of a cauliflower mosaic virus 35S promoter, a cassava vein mosaic virus promoter, and a cauliflower mosaic virus 35S minimal promoter, an elongation factor 1 alpha (EF1) promoter, a phosphoglycerate kinase (PGK) promoter, a ubiquitin (Ubc) promoter, and an albumin  
5 promoter. In addition, these expression sequences may be modified by addition of enhancer or regulatory sequences and the like.

Enhancers that may be used in embodiments of the invention include but are not limited to: tobacco mosaic virus enhancer, cauliflower mosaic virus 35S enhancer, tobacco etch virus enhancer, ribulose 1, 5-bisphosphate carboxylase enhancer, rice tungro bacilliform virus  
10 enhancer, and other plant and viral gene enhancers, and the like.

Termination control regions, *i.e.*, terminator or polyadenylation sequences, may also be derived from various genes native to the preferred hosts. Optionally, a termination site may be unnecessary, however, it is most preferred if included. In a preferred embodiment of the invention, the termination control region may be comprise or be derived from a synthetic  
15 sequence, synthetic polyadenylation signal, an SV40 late polyadenylation signal, an SV40 polyadenylation signal, a bovine growth hormone (BGH) polyadenylation signal, nopaline synthase (nos), cauliflower mosaic virus (CaMV), octopine synthase (ocs), Agrobacterium, viral, and plant terminator sequences, or the like.

The terms "3' non-coding sequences" or "3' untranslated region (UTR)" refer to DNA  
20 sequences located downstream (3') of a coding sequence and may comprise polyadenylation [poly(A)] recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor.


25 "Regulatory region" means a nucleic acid sequence which regulates the expression of a second nucleic acid sequence. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region) or may include sequences of a different origin that are responsible for expressing different proteins or even synthetic proteins (a heterologous region). In particular, the sequences can be sequences of  
30 prokaryotic, eukaryotic, or viral genes or derived sequences that stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, promoters, enhancers, transcriptional termination sequences, and signal sequences which direct the

polypeptide into the secretory pathways of the target cell.

A regulatory region from a "heterologous source" is a regulatory region that is not naturally associated with the expressed nucleic acid. Included among the heterologous regulatory regions are regulatory regions from a different species, regulatory regions from a different gene, hybrid regulatory sequences, and regulatory sequences which do not occur in nature, but which are designed by one having ordinary skill in the art.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. "Antisense RNA" refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene. The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that is not translated yet has an effect on cellular processes.

A "polypeptide" is a polymeric compound comprised of covalently linked amino acid residues. Amino acids have the following general structure:



Amino acids are classified into seven groups on the basis of the side chain R: (1) aliphatic side chains, (2) side chains containing a hydroxylic (OH) group, (3) side chains containing sulfur atoms, (4) side chains containing an acidic or amide group, (5) side chains containing a basic group, (6) side chains containing an aromatic ring, and (7) proline, an imino acid in which the side chain is fused to the amino group. A polypeptide of the invention preferably comprises at least about 14 amino acids.

A "protein" is a polypeptide that performs a structural or functional role in a living

cell.

An "isolated polypeptide" or "isolated protein" is a polypeptide or protein that is substantially free of those compounds that are normally associated therewith in its natural state (e.g., other proteins or polypeptides, nucleic acids, carbohydrates, lipids). "Isolated" is not  
5 meant to exclude artificial or synthetic mixtures with other compounds, or the presence of impurities which do not interfere with biological activity, and which may be present, for example, due to incomplete purification, addition of stabilizers, or compounding into a pharmaceutically acceptable preparation.

"Fragment" of a polypeptide according to the invention will be understood to mean a  
10 polypeptide whose amino acid sequence is shorter than that of the reference polypeptide and which comprises, over the entire portion with these reference polypeptides, an identical amino acid sequence. Such fragments may, where appropriate, be included in a larger polypeptide of which they are a part. Such fragments of a polypeptide according to the invention may have a length of 10, 15, 20, 30 to 40, 50, 100, 200 or 300 amino acids.

15 A "variant" of a polypeptide or protein is any analogue, fragment, derivative, or mutant which is derived from a polypeptide or protein and which retains at least one biological property of the polypeptide or protein. Different variants of the polypeptide or protein may exist in nature. These variants may be allelic variations characterized by differences in the nucleotide sequences of the structural gene coding for the protein, or may involve differential  
20 splicing or post-translational modification. The skilled artisan can produce variants having single or multiple amino acid substitutions, deletions, additions, or replacements. These variants may include, *inter alia*: (a) variants in which one or more amino acid residues are substituted with conservative or non-conservative amino acids, (b) variants in which one or more amino acids are added to the polypeptide or protein, (c) variants in which one or more of  
25 the amino acids includes a substituent group, and (d) variants in which the polypeptide or protein is fused with another polypeptide such as serum albumin. The techniques for obtaining these variants, including genetic ( suppressions, deletions, mutations, etc.), chemical, and enzymatic techniques, are known to persons having ordinary skill in the art. A variant polypeptide preferably comprises at least about 14 amino acids.

30 A "heterologous protein" refers to a protein not naturally produced in the cell.

A "mature protein" refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with

pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

The term "signal peptide" refers to an amino terminal polypeptide preceding the secreted mature protein. The signal peptide is cleaved from and is therefore not present in the mature protein. Signal peptides have the function of directing and translocating secreted proteins across cell membranes. Signal peptide is also referred to as signal protein.

A "signal sequence" is included at the beginning of the coding sequence of a protein to be expressed on the surface of a cell. This sequence encodes a signal peptide, N-terminal to the mature polypeptide, that directs the host cell to translocate the polypeptide. The term "translocation signal sequence" is used herein to refer to this sort of signal sequence. Translocation signal sequences can be found associated with a variety of proteins native to eukaryotes and prokaryotes, and are often functional in both types of organisms.

The term "homology" refers to the percent of identity between two polynucleotide or two polypeptide moieties. The correspondence between the sequence from one moiety to another can be determined by techniques known to the art. For example, homology can be determined by a direct comparison of the sequence information between two polypeptide molecules by aligning the sequence information and using readily available computer programs. Alternatively, homology can be determined by hybridization of polynucleotides under conditions that form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s) and size determination of the digested fragments.

As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (*e.g.*, the immunoglobulin superfamily) and homologous proteins from different species (*e.g.*, myosin light chain, etc.) (Reeck et al., 1987, Cell 50:667.). Such proteins (and their encoding genes) have sequence homology, as reflected by their high degree of sequence similarity.

Accordingly, the term "sequence similarity" in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that may or may not share a common evolutionary origin (*see* Reeck et al., 1987, Cell 50:667). As used herein, the term "homologous" in all its grammatical forms and spelling variations refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies and homologous proteins from different species (Reeck *et al.*, *supra*). Such proteins (and their encoding genes) have sequence homology, as reflected by



their high degree of sequence similarity. However, in common usage and in the instant application, the term "homologous," when modified with an adverb such as "highly," may refer to sequence similarity and not a common evolutionary origin.

In a specific embodiment, two DNA sequences are "substantially homologous" or  
5 "substantially similar" when at least about 50% (preferably at least about 75%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular  
10 system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook *et al.*, 1989, *supra*.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the protein encoded by the DNA sequence.  
15 "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotide bases that do not substantially affect the functional  
20 properties of the resulting transcript. It is therefore understood that the invention encompasses more than the specific exemplary sequences. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products.

Moreover, the skilled artisan recognizes that substantially similar sequences  
25 encompassed by this invention are also defined by their ability to hybridize, under stringent conditions (0.1X SSC, 0.1% SDS, 65°C and washed with 2X SSC, 0.1% SDS followed by 0.1X SSC, 0.1% SDS), with the sequences exemplified herein. Substantially similar nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 70% identical to the DNA sequence of the nucleic acid fragments reported herein.  
30 Preferred substantially nucleic acid fragments of the instant invention are those nucleic acid fragments whose DNA sequences are at least 80% identical to the DNA sequence of the nucleic acid fragments reported herein. More preferred nucleic acid fragments are at least 90% identical to the DNA sequence of the nucleic acid fragments reported herein. Even more

preferred are nucleic acid fragments that are at least 95% identical to the DNA sequence of the nucleic acid fragments reported herein.

Two amino acid sequences are "substantially homologous" or "substantially similar" when greater than about 40% of the amino acids are identical, or greater than 60% are similar (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, *Version 7*, Madison, Wisconsin) pileup program.

The term "corresponding to" is used herein to refer to similar or homologous sequences, whether the exact position is identical or different from the molecule to which the similarity or homology is measured. A nucleic acid or amino acid sequence alignment may include spaces. Thus, the term "corresponding to" refers to the sequence similarity, and not the numbering of the amino acid residues or nucleotide bases.

A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to putatively identify that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). In general, a sequence of ten or more contiguous amino acids or thirty or more nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene specific oligonucleotide probes comprising 20-30 contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12-15 bases may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises enough of the sequence to specifically identify and/or isolate a nucleic acid fragment comprising the sequence.

The term "percent identity", as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by

known methods, including but not limited to those described in: *Computational Molecular Biology* (Lesk, A. M., ed.) Oxford University Press, New York (1988); *Biocomputing: Informatics and Genome Projects* (Smith, D. W., ed.) Academic Press, New York (1993); *Computer Analysis of Sequence Data, Part I* (Griffin, A. M., and Griffin, H. G., eds.) Humana Press, New Jersey (1994); *Sequence Analysis in Molecular Biology* (von Heinje, G., ed.) Academic Press (1987); and *Sequence Analysis Primer* (Gribskov, M. and Devereux, J., eds.) Stockton Press, New York (1991). Preferred methods to determine identity are designed to give the best match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Sequence alignments and percent identity calculations may be performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences may be performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method may be selected: KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

The term "sequence analysis software" refers to any computer algorithm or software program that is useful for the analysis of nucleotide or amino acid sequences. "Sequence analysis software" may be commercially available or independently developed. Typical sequence analysis software will include but is not limited to the GCG suite of programs (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI), BLASTP, BLASTN, BLASTX (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), and DNASTAR (DNASTAR, Inc. 1228 S. Park St. Madison, WI 53715 USA). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the "default values" of the program referenced, unless otherwise specified. As used herein "default values" will mean any set of values or parameters which originally load with the software when first initialized.

"Synthetic genes" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form gene segments that are then enzymatically assembled to construct the entire gene. "Chemically synthesized", as related to a sequence of DNA, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of DNA may be accomplished using well established procedures, or automated chemical synthesis can

be performed using one of a number of commercially available machines. Accordingly, the genes can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host.

- 5 Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

### GENE EXPRESSION MODULATION SYSTEM OF THE INVENTION

- Applicants have now shown that separating the transactivation and DNA binding domains by placing them on two different proteins results in greatly reduced background activity in the absence of a ligand and significantly increased activity over background in the presence of a ligand. Applicants' improved gene expression system comprises two chimeric gene expression; the first encoding a DNA binding domain fused to a nuclear receptor polypeptide and the second encoding a transactivation domain fused to a nuclear receptor polypeptide. The interaction of the first protein with the second protein effectively tethers the DNA binding domain to the transactivation domain. Since the DNA binding and transactivation domains reside on two different molecules, the background activity in the absence of ligand is greatly reduced.

- In general, the inducible gene expression modulation system of the invention comprises
- 20 a) a first chimeric gene that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first hybrid polypeptide comprising i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor; and b) a second chimeric gene that is capable of being expressed in the host
- 25 cell comprising a polynucleotide sequence that encodes a second hybrid polypeptide comprising: i) a transactivation domain; and ii) a ligand binding domain comprising the ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the transactivation domain are from other than EcR, RXR, or USP; and wherein the ligand binding domains from the first hybrid polypeptide and the second hybrid polypeptide are different and
- 30 dimerize.

This two-hybrid system exploits the ability of a pair of interacting proteins to bring the transcription activation domain into a more favorable position relative to the DNA binding domain such that when the DNA binding domain binds to the DNA binding site on the gene,

the transactivation domain more effectively activates the promoter (see, for example, U.S. Patent No. 5,283,173). This two-hybrid system is a significantly improved inducible gene expression modulation system compared to the two systems disclosed in International Patent Applications PCT/US97/05330 and PCT/US98/14215.

5       The ecdysone receptor-based gene expression modulation system of the invention may be either heterodimeric and homodimeric. A functional EcR complex generally refers to a heterodimeric protein complex consisting of two members of the steroid receptor family, an ecdysone receptor protein obtained from various insects, and an ultraspiracle (USP) protein or the vertebrate homolog of USP, retinoid X receptor protein (see Yao, et al. (1993) Nature 366, 10 476-479; Yao, et al., (1992) Cell 71, 63-72). However, the complex may also be a homodimer as detailed below. The functional ecdysteroid receptor complex may also include additional protein(s) such as immunophilins. Additional members of the steroid receptor family of proteins, known as transcriptional factors (such as DHR38 or *betaFTZ-1*), may also be ligand dependent or independent partners for EcR, USP, and/or RXR. Additionally, other cofactors 15 may be required such as proteins generally known as coactivators (also termed adapters or mediators). These proteins do not bind sequence-specifically to DNA and are not involved in basal transcription. They may exert their effect on transcription activation through various mechanisms, including stimulation of DNA-binding of activators, by affecting chromatin structure, or by mediating activator-initiation complex interactions. Examples of such 20 coactivators include RIP140, TIF1, RAP46/Bag-1, ARA70, SRC-1/NCoA-1, TIF2/GRIP/NCoA-2, ACTR/AIB1/RAC3/pCIP as well as the promiscuous coactivator C response element B binding protein, CBP/p300 (for review see Glass et al, Curr. Opin. Cell Biol. 9:222-232, 1997). Also, protein cofactors generally known as corepressors (also known as repressors, silencers, or silencing mediators) may be required to effectively inhibit 25 transcriptional activation in the absence of ligand. These corepressors may interact with the unliganded ecdysone receptor to silence the activity at the response element. Current evidence suggests that binding of ligand changes the conformation of the receptor, which results in release of the corepressor and recruitment of the above described coactivators, thereby abolishing their silencing activity. Examples of corepressors include N-CoR and SMRT (for 30 review, see Horwitz et al. Mol Endocrinol. 10: 1167-1177, 1996). These cofactors may either be endogenous within the cell or organism, or may be added exogenously as transgenes to be expressed in either a regulated or unregulated fashion. Homodimer complexes of the ecdysone receptor protein, USP, or RXR may also be functional under some circumstances.

The ecdysone receptor complex typically includes proteins which are members of the nuclear receptor superfamily wherein all members are characterized by the presence of an amino-terminal transactivation domain, a DNA binding domain ("DBD"), and a ligand binding domain ("LBD") separated from the DBD by a hinge region. As used herein, the term "DNA binding domain" comprises a minimal peptide sequence of a DNA binding protein, up to the entire length of a DNA binding protein, so long as the DNA binding domain functions to associate with a particular response element. Members of the nuclear receptor superfamily are also characterized by the presence of four or five domains: A/B, C, D, E, and in some members F (see Evans, *Science* 240:889-895 (1988)). The "A/B" domain corresponds to the transactivation domain, "C" corresponds to the DNA binding domain, "D" corresponds to the hinge region, and "E" corresponds to the ligand binding domain. Some members of the family may also have another transactivation domain on the carboxy-terminal side of the LBD corresponding to "F".

The DBD is characterized by the presence of two cysteine zinc fingers between which are two amino acid motifs, the P-box and the D-box, which confer specificity for ecdysone response elements. These domains may be either native, modified, or chimeras of different domains of heterologous receptor proteins. This EcR receptor, like a subset of the steroid receptor family, also possesses less well defined regions responsible for heterodimerization properties. Because the domains of EcR, USP, and RXR are modular in nature, the LBD, DBD, and transactivation domains may be interchanged.

Gene switch systems are known that incorporate components from the ecdysone receptor complex. However, in these known systems, whenever EcR is used it is associated with native or modified DNA binding domains and transactivation domains on the same molecule. USP or RXR are typically used as silent partners. We have now shown that when DNA binding domains and transactivation domains are on the same molecule the background activity in the absence of ligand is high and that such activity is dramatically reduced when DNA binding domains and transactivation domains are on different molecules, that is, on each of two partners of a heterodimeric or homodimeric complex. This two-hybrid system also provides improved sensitivity to non-steroidal ligands for example, diacylhydrazines, when compared to steroidal ligands for example, ponasterone A ("PonA") or muristerone A ("MurA"). That is, when compared to steroids, the non-steroidal ligands provide higher activity at a lower concentration. In addition, since transactivation based on EcR gene switches is often cell-line dependent, it is easier to tailor switching system to obtain maximum

transactivation capability for each application. Furthermore, this two-hybrid system avoids some side effects due to overexpression of RXR that often occur when unmodified RXR is used as a switching partner. In this two-hybrid system, native DNA binding and transactivation domains of EcR or RXR are eliminated. As a result, these chimeric molecules  
5 have less chance of interacting with other steroid hormone receptors present in the cell resulting in reduced side effects.

Specifically, Applicants' invention relates to a gene expression modulation system comprising: a) a first gene expression cassette that is capable of being expressed in a host cell, wherein the first gene expression cassette comprises a polynucleotide that encodes a first  
10 polypeptide comprising i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor; and b) a second gene expression cassette that is capable of being expressed in the host cell, wherein the second gene expression cassette comprises a polynucleotide sequence that encodes a second polypeptide comprising i) a  
15 transactivation domain; and ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than ultraspiracle (USP); wherein the DNA binding domain and the transactivation domain are from other than EcR, RXR, or USP; wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

The present invention also relates to a gene expression modulation system according to  
20 the present invention further comprising c) a third gene expression cassette comprising: i) the response element to which the DNA-binding domain of the first polypeptide binds; ii) a promoter that is activated by the transactivation domain of the second polypeptide; and iii) the gene whose expression is to be modulated.

In a specific embodiment, the gene whose expression is to be modulated is a  
25 homologous gene with respect to the host cell. In another specific embodiment, the gene whose expression is to be modulated is a heterologous gene with respect to the host cell.

In a specific embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain.

In another specific embodiment, the ligand binding domain of the first polypeptide  
30 comprises a retinoid X receptor ligand binding domain.

In a specific embodiment, the ligand binding domain of the second polypeptide comprises an ecdysone receptor ligand binding domain.

In another specific embodiment, the ligand binding domain of the second polypeptide

comprises a retinoid X receptor ligand binding domain.

In a preferred embodiment, the ligand binding domain of the first polypeptide comprises an ecdysone receptor ligand binding domain, and the ligand binding domain of the second polypeptide comprises a retinoid X receptor ligand binding domain.

- 5 In another preferred embodiment, the ligand binding domain of the first polypeptide is from a retinoid X receptor polypeptide, and the ligand binding domain of the second polypeptide is from an ecdysone receptor polypeptide.

Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or  
 10 modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR. A truncation mutation may be made by any method used in the art, including but not limited to restriction endonuclease digestion/deletion, PCR-mediated/oligonucleotide-directed deletion, chemical mutagenesis, UV strand breakage, and the like.

- 15 Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothis virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly  
 20 *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm  
 25 (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

- Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55,  
 30 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a



specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the ecdysone receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR $_{\alpha}$ , RXR $_{\beta}$ , or RXR $_{\gamma}$  isoform.

Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ

ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the retinoid X receptor ligand binding domain comprises a polypeptide sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide)].

The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- $\kappa$ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is obtained from a VP16, GAL4, or NF- $\kappa$ B. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

The response element ("RE") may be any response element with a known DNA binding domain, or an analog, combination, or modification thereof. Preferably, the RE is an RE from GAL4 ("GAL4RE"), LexA, a steroid/thyroid hormone nuclear receptor RE, or a synthetic RE that recognizes a synthetic DNA binding domain. More preferably, the RE is a GAL4RE comprising a polynucleotide sequence of SEQ ID NO: 47 or a LexA 8X operon comprising a polynucleotide sequence of SEQ ID NO: 48. Preferably, the first hybrid protein is substantially free of a transactivation domain and the second hybrid protein is substantially free of a DNA binding domain. For purposes of this invention, "substantially free" means that the protein in question does not contain a sufficient sequence of the domain in question to provide activation or binding activity.

The ligands for use in the present invention as described below, when combined with the ligand binding domain of an EcR, USP, RXR, or another polypeptide which in turn are bound to the response element linked to a gene, provide the means for external temporal

regulation of expression of the gene. The binding mechanism or the order in which the various components of this invention bind to each other, that is, ligand to receptor, first polypeptide to response element, second polypeptide to promoter, etc., is not critical. Binding of the ligand to the ligand binding domains of an EcR, USP, RXR, or another protein, enables expression or  
5 suppression of the gene. This mechanism does not exclude the potential for ligand binding to EcR, USP, or RXR, and the resulting formation of active homodimer complexes (e.g. EcR+EcR or USP+USP). Preferably, one or more of the receptor domains can be varied producing a chimeric gene switch. Typically, one or more of the three domains, DBD, LBD, and transactivation domain, may be chosen from a source different than the source of the other  
10 domains so that the chimeric genes and the resulting hybrid proteins are optimized in the chosen host cell or organism for transactivating activity, complementary binding of the ligand, and recognition of a specific response element. In addition, the response element itself can be modified or substituted with response elements for other DNA binding protein domains such as the GAL-4 protein from yeast (see Sadowski, et al. (1988) *Nature*, 335:563-564) or LexA  
15 protein from *E. coli* (see Brent and Ptashne (1985), *Cell*, 43:729-736), or synthetic response elements specific for targeted interactions with proteins designed, modified, and selected for such specific interactions (see, for example, Kim, et al. (1997), *Proc. Natl. Acad. Sci., USA*, 94:3616-3620) to accommodate chimeric receptors. Another advantage of chimeric systems is that they allow choice of a promoter used to drive the gene expression according to a desired  
20 end result. Such double control can be particularly important in areas of gene therapy, especially when cytotoxic proteins are produced, because both the timing of expression as well as the cells wherein expression occurs can be controlled. When genes, operatively linked to a suitable promoter, are introduced into the cells of the subject, expression of the exogenous genes is controlled by the presence of the system of this invention. Promoters may be  
25 constitutively or inducibly regulated or may be tissue-specific (that is, expressed only in a particular type of cells) or specific to certain developmental stages of the organism.

#### GENE EXPRESSION CASSETTES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention  
30 comprises a novel gene expression cassette that is capable of being expressed in a host cell, wherein the gene expression cassette comprises a polynucleotide encoding a hybrid polypeptide. Thus, Applicants' invention also provides novel gene expression cassettes for use in the gene expression system of the invention.

Specifically, the present invention provides a gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide. The hybrid polypeptide comprises either 1) a DNA-binding domain that recognizes a response element and a ligand binding domain of a nuclear receptor or 2) a transactivation domain and a ligand binding domain of a nuclear receptor, wherein the transactivation domain is from a nuclear receptor other than an EcR, an RXR, or a USP.

In a specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain that recognizes a response element and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor.

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain that recognizes a response element and a retinoid X receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than a retinoid X receptor.

The DNA binding domain can be any DNA binding domain with a known response element, including synthetic and chimeric DNA binding domains, or analogs, combinations, or modifications thereof. Preferably, the DBD is a GAL4 DBD, a LexA DBD, a transcription factor DBD, a steroid/thyroid hormone nuclear receptor superfamily member DBD, a bacterial LacZ DBD, or a yeast put DBD. More preferably, the DBD is a GAL4 DBD [SEQ ID NO: 41 (polynucleotide) or SEQ ID NO: 42 (polypeptide)] or a LexA DBD [(SEQ ID NO: 43 (polynucleotide) or SEQ ID NO: 44 (polypeptide))].

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor.

In another specific embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain and a retinoid X receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

The transactivation domain (abbreviated "AD" or "TA") may be any steroid/thyroid hormone nuclear receptor AD, synthetic or chimeric AD, polyglutamine AD, basic or acidic amino acid AD, a VP16 AD, a GAL4 AD, an NF- $\kappa$ B AD, a BP64 AD, or an analog, combination, or modification thereof. Preferably, the AD is a synthetic or chimeric AD, or is

obtained from a VP16, GAL4, or NF-kB. Most preferably, the AD is a VP16 AD [SEQ ID NO: 45 (polynucleotide) or SEQ ID NO: 46 (polypeptide)].

Preferably, the ligand binding domain is an EcR or RXR related steroid/thyroid hormone nuclear receptor family member ligand binding domain, or analogs, combinations, or  
5 modifications thereof. More preferably, the LBD is from EcR or RXR. Even more preferably, the LBD is from a truncated EcR or RXR.

Preferably, the EcR is an insect EcR selected from the group consisting of a Lepidopteran EcR, a Dipteran EcR, an Arthropod EcR, a Homopteran EcR and a Hemipteran EcR. More preferably, the EcR for use is a spruce budworm *Choristoneura fumiferana* EcR  
10 ("CfEcR"), a *Tenebrio molitor* EcR ("TmEcR"), a *Manduca sexta* EcR ("MsEcR"), a *Heliothis virescens* EcR ("HvEcR"), a silk moth *Bombyx mori* EcR ("BmEcR"), a fruit fly *Drosophila melanogaster* EcR ("DmEcR"), a mosquito *Aedes aegypti* EcR ("AaEcR"), a blowfly *Lucilia capitata* EcR ("LcEcR"), a Mediterranean fruit fly *Ceratitis capitata* EcR ("CcEcR"), a locust *Locusta migratoria* EcR ("LmEcR"), an aphid *Myzus persicae* EcR  
15 ("MpEcR"), a fiddler crab *Uca pugilator* EcR ("UpEcR"), or an ixodid tick *Amblyomma americanum* EcR ("AmaEcR"). Even more preferably, the LBD is from spruce budworm (*Choristoneura fumiferana*) EcR ("CfEcR") or fruit fly *Drosophila melanogaster* EcR ("DmEcR").

Preferably, the LBD is from a truncated insect EcR. The insect EcR polypeptide  
20 truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the insect EcR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the insect  
25 EcR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the insect EcR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-  
30 domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the ecdysone receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ

ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the ecdysone receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

Preferably, the RXR polypeptide is a mouse *Mus musculus* RXR ("MmRXR") or a human *Homo sapiens* RXR ("HsRXR"). The RXR polypeptide may be an RXR $\alpha$ , RXR $\beta$ , or RXR $\gamma$  isoform.

10 Preferably, the LBD is from a truncated RXR. The RXR polypeptide truncation comprises a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the RXR polypeptide truncation comprises a  
15 deletion of at least a partial polypeptide domain. Even more preferably, the RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains  
20 deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the retinoid X receptor ligand binding domain is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of  
25 SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the retinoid X receptor ligand binding domain comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID  
30 NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

In a preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA

DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

5 In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ  
10 ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or  
15 a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the gene expression cassette encodes a hybrid  
20 polypeptide comprising a DNA-binding domain comprising a polypeptide sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID  
25 NO: 39, and SEQ ID NO: 40.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group  
30 consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ

ID NO: 46 and an ecdysone receptor ligand binding domain comprising a polypeptide sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

5 In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID  
10 NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

In another preferred embodiment, the gene expression cassette encodes a hybrid polypeptide comprising a transactivation domain comprising a polypeptide sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence  
15 selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

For purposes of this invention EcR and RXR also include synthetic and chimeric EcR and RXR and their homologs.

20

#### POLYNUCLEOTIDES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention comprises a gene expression cassette comprising a polynucleotide that encodes a truncated EcR or RXR polypeptide comprising a truncation mutation and is useful in methods of  
25 modulating the expression of a gene within a host cell.

Thus, the present invention also relates to a polynucleotide that encodes an EcR or RXR polypeptide comprising a truncation mutation. Specifically, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or ligand sensitivity.

30 Preferably, the truncation mutation results in a polynucleotide that encodes a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210,



215, 220, 225, 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR

5 polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

10 In a specific embodiment, the EcR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10. In a specific embodiment, the polynucleotide according to the invention encodes a ecdysone receptor polypeptide comprising an amino acid  
15 sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF, which comprises the last 33 carboxy-terminal amino acids of C domain), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-  
20 DE).

In another specific embodiment, the RXR polynucleotide according to the invention comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30. In another specific  
25 embodiment, the polynucleotide according to the invention encodes a truncated RXR polypeptide comprising an amino acid sequence consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-  
30 truncated EF), and SEQ ID NO: 40 (HsRXR-E).

In particular, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific

embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a  
5 truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that  
10 reduces non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF).

15 The present invention also relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation, wherein the mutation enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of  
20 the EcR or RXR polypeptide. In another specific embodiment, the present invention relates to an isolated polynucleotide encoding an EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated polynucleotide encoding an EcR polypeptide comprising a truncation mutation that enhances  
25 non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the polynucleotide comprises a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF).

The present invention also relates to an isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a  
30 heterodimer comprising the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises a polynucleotide sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF),

SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-truncated EF). In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have  
5 been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

### POLYPEPTIDES OF THE INVENTION

The novel ecdysone receptor-based inducible gene expression system of the invention  
10 comprises a polynucleotide that encodes a truncated EcR or RXR polypeptide and is useful in methods of modulating the expression of a gene within a host cell. Thus, the present invention also relates to an isolated truncated EcR or RXR polypeptide encoded by a polynucleotide or a gene expression cassette according to the invention. Specifically, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that affects  
15 ligand binding activity or ligand sensitivity encoded by a polynucleotide according to the invention.

The present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation. Specifically, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that affects ligand binding activity or  
20 ligand sensitivity.

Preferably, the truncation mutation results in a truncated EcR polypeptide or a truncated RXR polypeptide comprising a deletion of at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145, 150, 155, 160, 165, 170, 175, 180, 185, 190, 195, 200, 205, 210, 215, 220, 225,  
25 230, 235, 240, 245, 250, 255, 260, or 265 amino acids. More preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least a partial polypeptide domain. Even more preferably, the EcR or RXR polypeptide truncation comprises a deletion of at least an entire polypeptide domain. In a specific embodiment, the EcR or RXR polypeptide truncation comprises a deletion of at least an A/B-domain deletion, a C-domain deletion, a D-domain  
30 deletion, an E-domain deletion, an F-domain deletion, an A/B/C-domains deletion, an A/B/1/2-C-domains deletion, an A/B/C/D-domains deletion, an A/B/C/D/F-domains deletion, an A/B/F-domains, and an A/B/C/F-domains deletion. A combination of several complete and/or partial domain deletions may also be performed.

In a preferred embodiment, the isolated truncated ecdysone receptor polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), and SEQ ID NO: 10 (DmEcR-DE). In another preferred embodiment, the isolated truncated ecdysone receptor polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), and SEQ ID NO: 20 (DmEcR-DE).

In a preferred embodiment, the isolated truncated RXR polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF) and SEQ ID NO: 30 (HsRXR-E). In another preferred embodiment, the isolated truncated RXR polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF), SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-

E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncatedEF), and SEQ ID NO: 30 (HsRXR-E).

Thus, the present invention relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces ligand binding activity or ligand sensitivity of  
5 the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE), SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-  
10 DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 38 (HsRXR-EF), SEQ ID NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

In a specific embodiment, the present invention relates to an isolated EcR or RXR  
15 polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-  
20 DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 8 (DmEcR-DEF), or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces  
25 steroid binding activity or steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 18 (DmEcR-DEF), or SEQ ID NO: 19 (DmEcR-EF).

In another specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or  
30 non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO:

4 (CfEcR-EF) or SEQ ID NO: 9 (DmEcR-EF). Accordingly, the present invention also relates to an isolated truncated EcR or RXR polypeptide comprising a truncation mutation that reduces non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising  
 5 a truncation mutation that reduces non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polypeptide comprises an amino acid sequence of SEQ ID NO: 14 (CfEcR-EF) or SEQ ID NO: 19 (DmEcR-EF).

In particular, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of  
 10 the EcR or RXR polypeptide, wherein the polypeptide is encoded by a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 1 (CfEcR-CDEF), SEQ ID NO: 2 (CfEcR-1/2CDEF), SEQ ID NO: 3 (CfEcR-DEF), SEQ ID NO: 4 (CfEcR-EF), SEQ ID NO: 5 (CfEcR-DE), SEQ ID NO: 6 (DmEcR-CDEF), SEQ ID NO: 7 (DmEcR-1/2CDEF), SEQ ID NO: 8 (DmEcR-DEF), SEQ ID NO: 9 (DmEcR-EF),  
 15 SEQ ID NO: 10 (DmEcR-DE), SEQ ID NO: 21 (MmRXR-CDEF), SEQ ID NO: 22 (MmRXR-DEF), SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 25 (MmRXR-E), SEQ ID NO: 26 (HsRXR-CDEF), SEQ ID NO: 27 (HsRXR-DEF), SEQ ID NO: 28 (HsRXR-EF), SEQ ID NO: 29 (HsRXR-truncated EF), and SEQ ID NO: 30 (HsRXR-E).

20 The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11 (CfEcR-CDEF), SEQ ID NO: 12 (CfEcR-1/2CDEF), SEQ ID NO: 13 (CfEcR-DEF), SEQ ID NO: 14 (CfEcR-EF), SEQ ID NO: 15 (CfEcR-DE),  
 25 SEQ ID NO: 16 (DmEcR-CDEF), SEQ ID NO: 17 (DmEcR-1/2CDEF), SEQ ID NO: 18 (DmEcR-DEF), SEQ ID NO: 19 (DmEcR-EF), SEQ ID NO: 20 (DmEcR-DE), SEQ ID NO: 31 (MmRXR-CDEF), SEQ ID NO: 32 (MmRXR-DEF), SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 35 (MmRXR-E), SEQ ID NO: 36 (HsRXR-CDEF), SEQ ID NO: 37 (HsRXR-DEF), SEQ ID NO: 39 (HsRXR-EF), SEQ ID  
 30 NO: 39 (HsRXR-truncatedEF), and SEQ ID NO: 40 (HsRXR-E).

The present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances ligand binding activity or ligand sensitivity of the EcR or RXR polypeptide. In a specific embodiment, the present invention relates to an isolated EcR or

RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide.

5 In another specific embodiment, the present invention relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein  
10 the EcR polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 3 (CfEcR-DEF) or SEQ ID NO: 8 (DmEcR-DEF). Accordingly, the present invention also relates to an isolated EcR or RXR polypeptide comprising a truncation mutation that enhances non-steroid binding activity or steroid sensitivity of the EcR or RXR polypeptide. In a preferred embodiment, the present invention relates to an isolated EcR polypeptide comprising  
15 a truncation mutation that enhances non-steroid binding activity or non-steroid sensitivity of the EcR polypeptide, wherein the EcR polynucleotide comprises an amino acid sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

The present invention also relates to an isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprising  
20 the mutated retinoid X receptor polypeptide and a dimerization partner. Preferably, the isolated retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 23 (MmRXR-EF), SEQ ID NO: 24 (MmRXR-truncatedEF), SEQ ID NO: 28 (HsRXR-EF), or SEQ ID NO: 29 (HsRXR-  
25 truncatedEF). More preferably, the isolated polynucleotide encoding a retinoid X receptor polypeptide comprising a truncation mutation that increases ligand sensitivity of a heterodimer comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 33 (MmRXR-EF), SEQ ID NO: 34 (MmRXR-truncatedEF), SEQ ID NO: 38 (HsRXR-EF), or SEQ ID NO: 39 (HsRXR-truncatedEF).

30 In a specific embodiment, the dimerization partner is an ecdysone receptor polypeptide. Preferably, the dimerization partner is a truncated EcR polypeptide. More preferably, the dimerization partner is an EcR polypeptide in which domains A/B/C have been deleted. Even more preferably, the dimerization partner is an EcR polypeptide comprising an amino acid

sequence of SEQ ID NO: 13 (CfEcR-DEF) or SEQ ID NO: 18 (DmEcR-DEF).

#### METHOD OF MODULATING GENE EXPRESSION OF THE INVENTION

Applicants' invention also relates to methods of modulating gene expression in a host  
5 cell using a gene expression modulation system according to the invention. Specifically,  
Applicants' invention provides a method of modulating the expression of a gene in a host cell  
comprising the steps of: a) introducing into the host cell a gene expression modulation system  
according to the invention; and b) introducing into the host cell a ligand that independently  
combines with the ligand binding domains of the first polypeptide and the second polypeptide  
10 of the gene expression modulation system; wherein the gene to be expressed is a component of  
a gene expression cassette comprising: i) a response element comprising a domain to which the  
DNA binding domain of the first polypeptide binds; ii) a promoter that is activated by the  
transactivation domain of the second polypeptide; and iii) a gene whose expression is to be  
modulated, whereby a complex is formed comprising the ligand, the first polypeptide of the  
15 gene expression modulation system and the second polypeptide of the gene expression  
modulation system, and whereby the complex modulates expression of the gene in the host cell.


Genes of interest for expression in a host cell using Applicants' methods may be  
endogenous genes or heterologous genes. Nucleic acid or amino acid sequence information for  
a desired gene or protein can be located in one of many public access databases, for example,  
20 GENBANK, EMBL, Swiss-Prot, and PIR, or in many biology related journal publications.  
Thus, those skilled in the art have access to nucleic acid sequence information for virtually all  
known genes. Such information can then be used to construct the desired constructs for the  
insertion of the gene of interest within the gene expression cassettes used in Applicants'  
methods described herein.

25 Examples of genes of interest for expression in a host cell using Applicants' methods  
include, but are not limited to: antigens produced in plants as vaccines, enzymes like alpha-  
amylase, phytase, glucanase, and xylanase, genes for resistance against insects, nematodes, fungi,  
bacteria, viruses, and abiotic stresses, nutraceuticals, pharmaceuticals, vitamins, genes for  
modifying amino acid content, herbicide resistance, cold, drought, and heat tolerance, industrial  
30 products, oils, protein, carbohydrates, antioxidants, male sterile plants, flowers, fuels, other  
output traits, genes encoding therapeutically desirable polypeptides or products, such as genes  
that can provide, modulate, alleviate, correct and/or restore polypeptides important in treating a  
condition, a disease, a disorder, a dysfunction, a genetic defect, and the like.



Acceptable ligands are any that modulate expression of the gene when binding of the DNA binding domain of the two hybrid system to the response element in the presence of the ligand results in activation or suppression of expression of the genes. Preferred ligands include ponasterone, muristerone A, N,N'-diacylhydrazines such as those disclosed in U. S. Patents  
 5 No. 6,013,836; 5,117,057; 5,530,028; and 5,378,726; dibenzoylalkyl cyanohydrazines such as those disclosed in European Application No. 461,809; N-alkyl-N,N'-diaroylhydrazines such as those disclosed in U. S. Patent No. 5,225,443; N-acyl-N-alkylcarbonylhydrazines such as those disclosed in European Application No. 234,994; N-aroyl-N-alkyl-N'-aroylhydrazines such as those described in U. S. Patent No. 4,985,461; each of which is incorporated herein by  
 10 reference and other similar materials including 3,5-di-tert-butyl-4-hydroxy-N-isobutylbenzamide, 8-O-acetylharpagide, and the like.

Preferably, the ligand for use in Applicants' method of modulating expression of gene is a compound of the formula:



15 wherein:

E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl,  
 20 CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or  
 25 joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to

form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

5 Applicants' invention provides for modulation of gene expression in prokaryotic and eukaryotic host cells. Thus, the present invention also relates to a method for modulating gene expression in a host cell selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Preferably, the host cell is a yeast cell, a plant cell, a murine cell, or a human cell.

10 Expression in transgenic host cells may be useful for the expression of various polypeptides of interest including but not limited to therapeutic polypeptides, pathway intermediates; for the modulation of pathways already existing in the host for the synthesis of new products heretofore not possible using the host; cell based assays; and the like. Additionally the gene products may be useful for conferring higher growth yields of the host  
15 or for enabling alternative growth mode to be utilized.

#### HOST CELLS AND NON-HUMAN ORGANISMS OF THE INVENTION

As described above, the gene expression modulation system of the present invention may be used to modulate gene expression in a host cell. Expression in transgenic host cells  
20 may be useful for the expression of various genes of interest. Thus, Applicants' invention also provides an isolated host cell comprising a gene expression system according to the invention. The present invention also provides an isolated host cell comprising a gene expression cassette according to the invention. Applicants' invention also provides an isolated host cell comprising a polynucleotide or polypeptide according to the invention. The isolated  
25 host cell may be either a prokaryotic or a eukaryotic host cell.

Preferably, the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell. Examples of preferred host cells include, but are not limited to, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Candida*, *Hansenula*, or bacterial species such as  
30 those in the genera *Synechocystis*, *Synechococcus*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Rhodococcus*, *Streptomyces*, *Escherichia*, *Pseudomonas*, *Methylobacter*, *Methylobacter*, *Alcaligenes*, *Synechocystis*, *Anabaena*, *Thiobacillus*, *Methanobacterium* and *Klebsiella*, plant, animal, and mammalian host cells. More preferably, the host cell is a yeast cell, a plant

cell, a murine cell, or a human cell.

In a specific embodiment, the host cell is a yeast cell selected from the group consisting of a *Saccharomyces*, a *Pichia*, and a *Candida* host cell.

In another specific embodiment, the host cell is a plant cell selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, bean, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat host cell.

10 In another specific embodiment, the host cell is a murine cell.

In another specific embodiment, the host cell is a human cell.

Host cell transformation is well known in the art and may be achieved by a variety of methods including but not limited to electroporation, viral infection, plasmid/vector transfection, non-viral vector mediated transfection, *Agrobacterium*-mediated transformation, particle bombardment, and the like. Expression of desired gene products involves culturing the transformed host cells under suitable conditions and inducing expression of the transformed gene. Culture conditions and gene expression protocols in prokaryotic and eukaryotic cells are well known in the art (see General Methods section of Examples). Cells may be harvested and the gene products isolated according to protocols specific for the gene product.

20 In addition, a host cell may be chosen which modulates the expression of the inserted polynucleotide, or modifies and processes the polypeptide product in the specific fashion desired. Different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (*e.g.*, glycosylation, cleavage [*e.g.*, of signal sequence]) of proteins. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system can be used to produce a non-glycosylated core protein product.

25 However, a polypeptide expressed in bacteria may not be properly folded. Expression in yeast can produce a glycosylated product. Expression in eukaryotic cells can increase the likelihood of "native" glycosylation and folding of a heterologous protein. Moreover, expression in mammalian cells can provide a tool for reconstituting, or constituting, the polypeptide's activity. Furthermore, different vector/host expression systems may affect processing reactions, such as proteolytic cleavages, to a different extent.

Applicants' invention also relates to a non-human organism comprising an isolated host cell according to the invention. Preferably, the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal. More preferably, the non-human organism is a yeast, a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, or a chimpanzee.

In a specific embodiment, the non-human organism is a yeast selected from the group consisting of *Saccharomyces*, *Pichia*, and *Candida*.

In another specific embodiment, the non-human organism is a plant selected from the group consisting of an apple, *Arabidopsis*, bajra, banana, barley, beans, beet, blackgram, chickpea, chili, cucumber, eggplant, favabean, maize, melon, millet, mungbean, oat, okra, *Panicum*, papaya, peanut, pea, pepper, pigeonpea, pineapple, *Phaseolus*, potato, pumpkin, rice, sorghum, soybean, squash, sugarcane, sugarbeet, sunflower, sweet potato, tea, tomato, tobacco, watermelon, and wheat.

In another specific embodiment, the non-human organism is a *Mus musculus* mouse.

#### MEASURING GENE EXPRESSION/TRANSCRIPTION

One useful measurement of Applicants' methods of the invention is that of the transcriptional state of the cell including the identities and abundances of RNA, preferably mRNA species. Such measurements are conveniently conducted by measuring cDNA abundances by any of several existing gene expression technologies.

Nucleic acid array technology is a useful technique for determining differential mRNA expression. Such technology includes, for example, oligonucleotide chips and DNA microarrays. These techniques rely on DNA fragments or oligonucleotides which correspond to different genes or cDNAs which are immobilized on a solid support and hybridized to probes prepared from total mRNA pools extracted from cells, tissues, or whole organisms and converted to cDNA. Oligonucleotide chips are arrays of oligonucleotides synthesized on a substrate using photolithographic techniques. Chips have been produced which can analyze for up to 1700 genes. DNA microarrays are arrays of DNA samples, typically PCR products, that are robotically printed onto a microscope slide. Each gene is analyzed by a full or partial-length target DNA sequence. Microarrays with up to 10,000 genes are now routinely prepared commercially. The primary difference between these two techniques is that oligonucleotide chips typically utilize 25-mer oligonucleotides which allow fractionation of short DNA molecules whereas the larger DNA targets of microarrays, approximately 1000 base pairs, may

provide more sensitivity in fractionating complex DNA mixtures.

Another useful measurement of Applicants' methods of the invention is that of determining the translation state of the cell by measuring the abundances of the constituent protein species present in the cell using processes well known in the art.

5       Where identification of genes associated with various physiological functions is desired, an assay may be employed in which changes in such functions as cell growth, apoptosis, senescence, differentiation, adhesion, binding to a specific molecules, binding to another cell, cellular organization, organogenesis, intracellular transport, transport facilitation, energy conversion, metabolism, myogenesis, neurogenesis, and/or hematopoiesis is measured.

10       In addition, selectable marker or reporter gene expression may be used to measure gene expression modulation using Applicants' invention.

Other methods to detect the products of gene expression are well known in the art and include Southern blots (DNA detection), dot or slot blots (DNA, RNA), Northern blots (RNA), and RT-PCR (RNA) analyses. Although less preferred, labeled proteins can be used to detect  
15 a particular nucleic acid sequence to which it hybridizes.

In some cases it is necessary to amplify the amount of a nucleic acid sequence. This may be carried out using one or more of a number of suitable methods including, for example, polymerase chain reaction ("PCR"), ligase chain reaction ("LCR"), strand displacement amplification ("SDA"), transcription-based amplification, and the like. PCR is carried out in  
20 accordance with known techniques in which, for example, a nucleic acid sample is treated in the presence of a heat stable DNA polymerase, under hybridizing conditions, with one oligonucleotide primer for each strand of the specific sequence to be detected. An extension product of each primer that is synthesized is complementary to each of the two nucleic acid strands, with the primers sufficiently complementary to each strand of the specific sequence to  
25 hybridize therewith. The extension product synthesized from each primer can also serve as a template for further synthesis of extension products using the same primers. Following a sufficient number of rounds of synthesis of extension products, the sample may be analyzed as described above to assess whether the sequence or sequences to be detected are present.

30       The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention.

## EXAMPLES

### GENERAL METHODS

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described by Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 5 (1989) (Maniatis) and by T. J. Silhavy, M. L. Bennis, and L. W. Enquist, *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1984) and by Ausubel, F. M. et al., *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley-Interscience (1987).

10 Methods for plant tissue culture, transformation, plant molecular biology, and plant, general molecular biology may be found in *Plant Tissue Culture Concepts and Laboratory Exercises* edited by RN Trigiano and DJ Gray, 2<sup>nd</sup> edition, 2000, CRC press, New York; *Agrobacterium Protocols* edited by KMA Gartland and MR Davey, 1995, Humana Press, Totowa, New Jersey; *Methods in Plant Molecular Biology*, P. Maliga et al., 1995, Cold 15 Spring Harbor Lab Press, New York; and *Molecular Cloning*, J. Sambrook et al., 1989, Cold Spring Harbor Lab Press, New York.

Materials and methods suitable for the maintenance and growth of bacterial cultures are well known in the art. Techniques suitable for use in the following examples may be found as set out in *Manual of Methods for General Bacteriology* (Phillipp Gerhardt, R. G. E. 20 Murray, Ralph N. Costilow, Eugene W. Nester, Willis A. Wood, Noel R. Krieg and G. Briggs Phillips, eds), American Society for Microbiology, Washington, DC. (1994)) or by Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition, Sinauer Associates, Inc., Sunderland, MA (1989). All reagents, restriction enzymes and materials used for the growth and maintenance of host cells were obtained from Aldrich Chemicals 25 (Milwaukee, WI), DIFCO Laboratories (Detroit, MI), GIBCO/BRL (Gaithersburg, MD), or Sigma Chemical Company (St. Louis, MO) unless otherwise specified.

Manipulations of genetic sequences may be accomplished using the suite of programs available from the Genetics Computer Group Inc. (Wisconsin Package Version 9.0, Genetics Computer Group (GCG), Madison, WI). Where the GCG program "Pileup" is used the gap 30 creation default value of 12, and the gap extension default value of 4 may be used. Where the CGC "Gap" or "Bestfit" programs is used the default gap creation penalty of 50 and the default gap extension penalty of 3 may be used. In any case where GCG program parameters are not prompted for, in these or any other GCG program, default values may be used.

The meaning of abbreviations is as follows: "h" means hour(s), "min" means minute(s), "sec" means second(s), "d" means day(s), "μl" means microliter(s), "ml" means milliliter(s), "L" means liter(s), "μM" means micromolar, "mM" means millimolar, "μg" means microgram(s), "mg" means milligram(s), "A" means adenine or adenosine, "T" means thymine or thymidine, "G" means guanine or guanosine, "C" means cytidine or cytosine, "x g" means times gravity, "nt" means nucleotide(s), "aa" means amino acid(s), "bp" means base pair(s), "kb" means kilobase(s), "k" means kilo, "μ" means micro, and "°C" means degrees Celsius.

10

### EXAMPLE 1

Applicants' improved EcR-based inducible gene modulation system was developed for use in various applications including gene therapy, expression of proteins of interest in host cells, production of transgenic organisms, and cell-based assays. This Example describes the construction and evaluation of several gene expression cassettes for use in the EcR-based inducible gene expression system of the invention.

In various cellular backgrounds, including mammalian cells, insect ecdysone receptor (EcR) heterodimerizes with retinoid X receptor (RXR) and, upon binding of ligand, transactivates genes under the control of ecdysone response elements. Applicants constructed several EcR-based gene expression cassettes based on the spruce budworm *Choristoneura fumiferana* EcR ("CfEcR"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 49 and SEQ ID NO: 50, respectively), *C. fumiferana* ultraspiracle ("CfUSP"; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 51 and SEQ ID NO: 52, respectively), and mouse *Mus musculus* RXRα (MmRXRα; full length polynucleotide and amino acid sequences are set forth in SEQ ID NO: 53 and SEQ ID NO: 54, respectively). The prepared receptor constructs comprise a ligand binding domain of EcR and of RXR or of USP; a DNA binding domain of GAL4 or of EcR; and an activation domain of VP16. The reporter constructs include a reporter gene, luciferase or LacZ, operably linked to a synthetic promoter construct that comprises either GAL4 or EcR/USP binding sites (response elements). Various combinations of these receptor and reporter constructs were cotransfected into CHO, NIH3T3, CV1 and 293 cells. Gene induction potential (magnitude of induction) and ligand specificity and sensitivity were examined using four different ligands: two steroidal ligands (ponasterone A and muristerone A) and two non-steroidal ligands (N-(2-ethyl-3-

methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine) in a dose-dependent induction of reporter gene expression in the transfected cells. Reporter gene expression activities were assayed at 24hr or 48hr after ligand addition.

5

**Gene Expression Cassettes:** Ecdysone receptor-based, chemically inducible gene expression cassettes (switches) were constructed as followed, using standard cloning methods available in the art. The following is brief description of preparation and composition of each switch.

1.1 - GAL4EcR/VP16RXR: The D, E, and F domains from spruce budworm *Choristoneura fumiferana* EcR ("CfEcRDEF"; SEQ ID NO: 3) were fused to GAL4 DNA binding domain ("DNABD"; SEQ ID NO: 41) and placed under the control of an SV40e promoter (SEQ ID NO: 55). The DEF domains from mouse (*Mus musculus*) RXR ("MmRXRDEF"; SEQ ID NO: 22) were fused to the activation domain from VP16 ("VP16AD"; SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Five consensus GAL4 binding sites ("5XGAL4RE"; comprising 5, GAL4RE comprising SEQ ID NO: 47) were fused to a synthetic E1b minimal promoter (SEQ ID NO: 56) and placed upstream of the luciferase gene (SEQ ID NO: 57).

1.2 - GAL4EcR/VP16USP: This construct was prepared in the same way as in switch 1.1 above except MmRXRDEF was replaced with the D, E and F domains from spruce budworm USP ("CfUSPDEF"; SEQ ID NO: 58). The constructs used in this example are similar to those disclosed in U. S. Patent No. 5,880,333 except that *Choristoneura fumiferana* USP rather than *Drosophila melanogaster* USP was utilized.

1.3 - GAL4RXR/VP16CfEcR: MmRXRDEF (SEQ ID NO: 22) was fused to a GAL4DNABD (SEQ ID NO: 41) and CfEcRCDEF (SEQ ID NO: 1) was fused to a VP16AD (SEQ ID NO: 45).

1.4 - GAL4RXR/VP16DmEcR: This construct was prepared in the same way as switch 1.3 except CfEcRCDEF was replaced with DmEcRCDEF (SEQ ID NO: 6).

1.5 - GAL4USP/VP16CfEcR: This construct was prepared in the same way as switch 1.3 except MmRXRDEF was replaced with CfUSPDEF (SEQ ID NO: 58).

1.6 - GAL4RXRCfEcRV16: This construct was prepared so that both the GAL4 DNABD and the VP16AD were placed on the same molecule. GAL4DNABD (SEQ ID NO: 41) and VP16AD (SEQ ID NO: 45) were fused to CfEcRDEF (SEQ ID NO: 3) at N-and C-termini respectively. The fusion was placed under the control of an SV40e promoter (SEQ ID NO:

30



55).

1.7 - VP16CfEcR: This construct was prepared such that CfEcRCDEF (SEQ ID NO: 1) was fused to VP16AD (SEQ ID NO: 45) and placed under the control of an SV40e promoter (SEQ ID NO: 55). Six ecdysone response elements ("EcRE"; SEQ ID NO: 59) from the hsp27 gene  
5 were placed upstream of the promoter and a luciferase gene (SEQ ID NO: 57). This switch most probably uses endogenous RXR.

1.8 - DmVgRXR: This system was purchased from Invitrogen Corp., Carlsbad, California. It comprises a *Drosophila melanogaster* EcR ("DmEcR") with a modified DNABD fused to VP16AD and placed under the control of a CMV promoter (SEQ ID NO: 60). Full length  
10 MmRXR (SEQ ID NO: 53) was placed under the control of the RSV promoter (SEQ ID NO: 61). The reporter, pIND(SP1)LacZ, contains five copies of a modified ecdysone response element ("EcRE", E/GRE), three copies of an SP1 enhancer, and a minimal heat shock promoter, all of which were placed upstream to the LacZ reporter gene.

1.9 - CfVgRXR: This example was prepared in the same way as switch 1.8 except DmEcR  
15 was replaced with a truncated CfEcR comprising a partial A/B domain and the complete CDEF domains [SEQ ID NO: 62 (polynucleotide) and SEQ ID NO: 63 (polypeptide)].

1.10 - CfVgRXRdel: This example was prepared in the same way as switch 1.9 except MmRXR (SEQ ID NO: 53) was deleted.

20 **Cell lines:** Four cell lines: CHO, Chinese hamster *Cricetulus griseus* ovarian cell line; NIH3T3 (3T3) mouse *Mus musculus* cell line; 293 human *Homo sapiens* kidney cell line, and CV1 African green monkey kidney cell line were used in these experiments. Cells were maintained in their respective media and were subcultured when they reached 60% confluency. Standard methods for culture and maintenance of the cells were followed.

25

**Transfections:** Several commercially available lipofactors as well as electroporation methods were evaluated and the best conditions for transfection of each cell line were developed. CHO, NIH3T3, 293 and CV1 cells were grown to 60% confluency. DNAs corresponding to the various switch constructs outlined in Examples 1.1 through 1.10 were transfected into CHO  
30 cells, NIH3T3 cells, 293 cells, or CV1 cells as follows.

CHO cells: Cells were harvested when they reach 60-80% confluency and plated in 6- or 12- or 24- well plates at 250,000, 100,000, or 50,000 cells in 2.5, 1.0, or 0.5 ml of growth medium containing 10% Fetal bovine serum respectively. The next day, the cells were rinsed with

growth medium and transfected for four hours. LipofectAMINE™ 2000 (Life Technologies Inc.) was found to be the best transfection reagent for these cells. For 12- well plates, 4 µl of LipofectAMINE™ 2000 was mixed with 100 µl of growth medium. 1.0 µg of reporter construct and 0.25 µg of receptor construct(s) were added to the transfection mix. A second  
5 reporter construct was added [pTKRL (Promega), 0.1 µg/transfection mix] and comprised a *Renilla* luciferase gene (SEQ ID NO: 64) operably linked and placed under the control of a thymidine kinase (TK) constitutive promoter and was used for normalization. The contents of the transfection mix were mixed in a vortex mixer and let stand at room temperature for 30 min. At the end of incubation, the transfection mix was added to the cells maintained in 400 µl  
10 growth medium. The cells were maintained at 37°C and 5% CO<sub>2</sub> for four hours. At the end of incubation, 500 µl of growth medium containing 20% FBS and either DMSO (control) or a DMSO solution of appropriate ligands were added and the cells were maintained at 37 °C and 5% CO<sub>2</sub> for 24-48 hr. The cells were harvested and reporter activity was assayed. The same procedure was followed for 6 and 24 well plates as well except all the reagents were doubled  
15 for 6 well plates and reduced to half for 24-well plates.

NIH3T3 Cells: Superfect™ (Qiagen Inc.) was found to be the best transfection reagent for 3T3 cells. The same procedures described for CHO cells were followed for 3T3 cells as well with two modifications. The cells were plated when they reached 50% confluency. 125,000 or 50,000 or 25,000 cells were plated per well of 6- or 12- or 24-well plates respectively. The  
20 GA14EcR/VP16RXR and reporter vector DNAs were transfected into NIH3T3 cells, the transfected cells were grown in medium containing PonA, MurA, N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine for 48 hr. The ligand treatments were performed as described in the CHO cell section above.

25 293 Cells: LipofectAMINE™ 2000 (Life Technologies) was found to be the best lipofactor for 293 cells. The same procedures described for CHO were followed for 293 cells except that the cells were plated in biocoated plates to avoid clumping. The ligand treatments were performed as described in the CHO cell section above.

CV1 Cells: LipofectAMINE™ plus (Life Technologies) was found to be the best lipofactor  
30 for CV1 cells. The same procedures described for NIH3T3 cells were followed for CV1 cells

**Ligands:** Ponasterone A and Muristerone A were purchased from Sigma Chemical Company. The two non-steroids N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-

butylhydrazine, or N-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine are synthetic stable ecdysteroids synthesized at Rohm and Haas Company. All ligands were dissolved in DMSO and the final concentration of DMSO was maintained at 0.1% in both controls and treatments.

5

**Reporter Assays:** Cells were harvested 24-48 hr after adding ligands. 125, 250, or 500  $\mu$ l of passive lysis buffer (part of Dual-luciferase<sup>TM</sup> reporter assay system from Promega Corporation) were added to each well of 24- or 12- or 24-well plates respectively. The plates were placed on a rotary shaker for 15 min. Twenty  $\mu$ l of lysate was assayed. Luciferase

- 10 activity was measured using Dual-luciferase<sup>TM</sup> reporter assay system from Promega Corporation following the manufacturer's instructions.  $\beta$ -Galactosidase was measured using Galacto-Star<sup>TM</sup> assay kit from TROPIX following the manufacturer's instructions. All luciferase and  $\beta$ -galactosidase activities were normalized using *Renilla* luciferase as a standard. Fold activities were calculated by dividing normalized relative light units ("RLU") in
- 15 ligand treated cells with normalized RLU in DMSO treated cells (untreated control).

The results of these experiments are provided in the following tables.

Table 1

## Transactivation of reporter genes through various switches in CHO cells

20

Composition of Switch	Mean Fold Activation with 50 $\mu$ M N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	267
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	85
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	312
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2
1.6 GAL4CfEcRVP16 pGAL4RELuc	9
1.7 VP16CfEcR pEcRELuc	36
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	14

1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	27
1.10 CfVgRXR pIND(SP1)LacZ	29

**Table 2****Transactivation of reporter genes through various switches in 3T3 cells**

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	1118
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	47
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	269
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3
1.6 GAL4CfEcRVP16 pGAL4RELuc	7
1.7 VP16CfEcR pEcRELuc	1
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	21
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	19
1.10 CfVgRXR pIND(SP1)LacZ	2

5

**Table 3****Transactivation of reporter genes through various switches in 293 cells**

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	125
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	17
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	3
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	2
1.6 GAL4CfEcRVP16 pGAL4RELuc	3

1.7 VP16CfEcR pEcRELuc	2
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	21
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	12
1.10 CfVgRXR pIND(SP1)LacZ	3

**Table 4**  
**Transactivation of reporter genes through various switches in CV1 cells**

Composition of Switch	Mean Fold Activation Through N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-t-butylhydrazine
1.1 GAL4EcR + VP16RXR pGAL4RELuc	279
1.2 GAL4EcR + VP16USP pGAL4RELuc	2
1.3 GAL4RXR + VP16CfEcR pGAL4RELuc	25
1.4 GAL4RXR + VP16DmEcR pGAL4RELuc	80
1.5 GAL4USP + VP16CfEcR pGAL4RELuc	3
1.6 GAL4CfEcRVP16 pGAL4RELuc	6
1.7 VP16CfEcR pEcRELuc	1
1.8 DmVgRXR + MmRXR pIND(SP1)LacZ	12
1.9 CfVgRXR + MmRXR pIND(SP1)LacZ	7
1.10 CfVgRXR pIND(SP1)LacZ	1

5

**Table 5**  
**Transactivation of reporter gene GAL4CfEcRDEF/VP16MmRXRDEF (switch 1.1)**  
**through steroids and non-steroids in 3T3 cells.**

Ligand	Mean Fold Induction at 1.0 $\mu$ M Concentration
1. Ponasterone A	1.0
2. Muristerone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	~ 116
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	601

10

**Table 6**  
**Transactivation of reporter gene GAL4MmRXRDEF/VP16CfEcRCDEF (switch 1.3)**  
**through steroids and non-steroids in 3T3 cells.**

Ligand	Mean Fold Induction at 1.0 $\mu$ M Concentration
1. Ponasterone A	1.0
2. Muristerone A	1.0
3. N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	71
4. N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine	54

5

Applicants' results demonstrate that the non-steroidal ecdysone agonists, N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine and N'-(3,4-(1,2-ethylenedioxy)-2-methylbenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine, were more potent activators of CfEcR as compared to *Drosophila melanogaster* EcR (DmEcR). (see  
 10 Tables 1-4). Also, in the mammalian cell lines tested, MmRXR performed better than CfUSP as a heterodimeric partner for CfEcR. (see Tables 1-4). Additionally, Applicants' inducible gene expression modulation system performed better when exogenous MmRXR was used than when the system relied only on endogenous RXR levels (see Tables 1-4).

Applicants' results also show that in a CfEcR-based inducible gene expression system,  
 15 the non-steroidal ecdysone agonists induced reporter gene expression at a lower concentration (i.e., increased ligand sensitivity) as compared to the steroid ligands, ponasterone A and muristerone A (see Tables 5 and 6).

Out of 10 EcR based gene switches tested, the GAL4EcR/VP16RXR switch (Switch 1.1) performed better than any other switch in all four cell lines examined and was more  
 20 sensitive to non-steroids than steroids. The results also demonstrate that placing the activation domain (AD) and DNA binding domain (DNABD) on each of the two partners reduced background when compared to placing both AD and DNABD together on one of the two partners. Therefore, a switch format where the AD and DNABD are separated between two partners, works well for EcR-based gene switch applications.

25 In addition, the MmRXR/EcR-based switches performed better than CfUSP/EcR-based switches, which have a higher background activity than the MmRXR/EcR switches in the absence of ligand.

Finally, the GAL4EcR/VP16RXR switch (Switch 1.1) was more sensitive to non-steroid ligands than to the steroid ligands (see Tables 5 and 6). In particular, steroid ligands

initiated transactivation at concentrations of 50  $\mu$ M, whereas the non-steroid ligands initiated transactivation at less than 1  $\mu$ M (submicromolar) concentration.

## EXAMPLE 2

5

This Example describes Applicants' further analysis of truncated EcR and RXR polypeptides in the improved EcR-based inducible gene expression system of the invention. To identify the best combination and length of two receptors that give a switch with a) maximum induction in the presence of ligand; b) minimum background in the absence of ligand; c) highly sensitive to ligand concentration; and d) minimum cross-talk among ligands and receptors, Applicants made and analyzed several truncation mutations of the CfEcR and MmRXR receptor polypeptides in NIH3T3 cells.

Briefly, polynucleotides encoding EcR or RXR receptors were truncated at the junctions of A/B, C, D, E and F domains and fused to either a GAL4 DNA binding domain encoding polynucleotide (SEQ ID NO: 41) for CfEcR, or a VP16 activation domain encoding polynucleotide (SEQ ID NO: 45) for MmRXR as described in Example 1. The resulting receptor truncation/fusion polypeptides were assayed in NIH3T3 cells. Plasmid pFRLUC (Stratagene) encoding a luciferase polypeptide was used as a reporter gene construct and pTKRL (Promega) encoding a *Renilla* luciferase polypeptide under the control of the constitutive TK promoter was used to normalize the transfections as described above. The analysis was performed in triplicates and mean luciferase counts were determined as described above.

### Gene Expression Cassettes Encoding Truncated Ecdysone Receptor Polypeptides

Gene expression cassettes comprising polynucleotides encoding either full length or truncated CfEcR polypeptides fused to a GAL4 DNA binding domain (SEQ ID NO: 41): GAL4CfEcRA/BCDEF (full length CfEcRA/BCDEF; SEQ ID NO: 49), GAL4CfEcRCDEF (CfEcRCDEF; SEQ ID NO: 1), GAL4CfEcR1/2CDEF (CfEcR1/2CDEF; SEQ ID NO: 2), GAL4CfEcRDEF (CfEcRDEF; SEQ ID NO: 3), GAL4CfEcREF (CfEcREF; SEQ ID NO: 4), and GAL4CfEcRDE (CfEcRDE; SEQ ID NO: 5) were transfected into NIH3T3 cells along with VP16MmRXRDEF (constructed as in Example 1.1; Figure 11) or VP16MmRXREF [constructed as in Example 1.1 except that MmRXRDEF was replaced with MmRXREF (SEQ ID NO: 23); Figure 12], and pFRLUC and pTKRL plasmid DNAs. The transfected cells were grown in the presence 0, 1, 5 or 25  $\mu$ M of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-

dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested, lysed and luciferase reporter activity was measured in the cell lysates. Total fly luciferase relative light units are presented. The number on the top of each bar is the maximum fold induction for that treatment.

- 5 Applicants' results show that the EF domain of MmRXR is sufficient and performs better than DEF domains of this receptor (see Figures 11 and 12). Applicants have also shown that, in general, EcR/RXR receptor combinations are insensitive to PonA (see Figures 11 and 12). As shown in the Figures 11 and 12, the GAL4CfEcRCDEF hybrid polypeptide (SEQ ID NO: 7) performed better than any other CfEcR hybrid polypeptide.

10 Gene Expression Cassettes Encoding Truncated Retinoid X Receptor Polypeptides

- Gene expression cassettes comprising polynucleotides encoding either full length or truncated MmRXR polypeptides fused to a VP16 transactivation domain (SEQ ID NO: 45): VP16MmRXRA/BCDEF (full length MmRXRA/BCDEF; SEQ ID NO: 53), VP16MmRXRCDEF (MmRXRCDEF; SEQ ID NO: 21), VP16MmRXRDEF (MmRXRDEF; SEQ ID NO: 22), VP16MmRXREF (MmRXREF; SEQ ID NO: 23), VP16MmRXRBam-EF ("MmRXRBam-EF" or "MmRXR-truncatedEF"; SEQ ID NO: 24), and VP16MmRXRAF2del ("MmRXRAF2del" or "MmRXR-E"; SEQ ID NO: 25) constructs were transfected into NIH3T3 cells along with GAL4CfEcRCDEF (constructed as in Example 1.1; Figure 13) or GAL4CfEcRDEF [constructed as in Example 1.1 except CfEcRCDEF was replaced with CfEcRDEF (SEQ ID NO: 3); Figure 14], pFRLuc and pTKRL plasmid DNAs as described above. The transfected cells were grown in the presence 0, 1, 5 and 25 uM of N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine or PonA for 48 hr. The cells were harvested and lysed and reporter activity was measured in the cell lysate. Total fly luciferase relative light units are presented. The number on top of each bar is the maximum fold induction in that treatment.

- Of all the truncations of MmRXR tested, Applicants' results show that the MmRXREF receptor was the best partner for CfEcR (Figures 13 and 14). CfEcRCDEF showed better induction than CfEcRDEF using MmRXREF. Deleting AF2 (abbreviated "EF-AF2del") or helices 1-3 of the E domain (abbreviated "EF-Bamdel") resulted in an RXR receptor that reduced gene induction and ligand sensitivity when partnered with either CfEcRCDEF (Figure 13) or CfEcRDEF (Figure 14) in NIH3T3 cells. In general, the CfEcR/RXR-based switch was much more sensitive to the non-steroid N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine than to the steroid PonA.



**EXAMPLE 3**

This Example describes Applicants' further analysis of gene expression cassettes encoding truncated EcR or RXR receptor polypeptides that affect either ligand binding activity or ligand sensitivity, or both. Briefly, six different combinations of chimeric receptor pairs, constructed as described in Examples 1 and 2, were further analyzed in a single experiment in NIH3T3 cells. These six receptor pair combinations and their corresponding sample numbers are depicted in Table 7.

10

**Table 7****CfEcR + MmRXR Truncation Receptor Combinations in NIH3T3 Cells**

<b>Figure 15 X-Axis Sample No.</b>	<b>EcR Polypeptide Construct</b>	<b>RXR Polypeptide Construct</b>
Samples 1 and 2	GAL4CfEcRCDEF	VP16RXRA/BCDEF (Full length)
Samples 3 and 4	GAL4CfEcRCDEF	VP16RXRDEF
Samples 5 and 6	GAL4CfEcRCDEF	VP16RXREF
Samples 7 and 8	GAL4CfEcRDEF	VP16RXRA/BCDEF (Full length)
Samples 9 and 10	GAL4CfEcRDEF	VP16RXRDEF
Samples 11 and 12	GAL4CfEcRDEF	VP16RXREF

The above receptor construct pairs, along with the reporter plasmid pFRLuc were transfected into NIH3T3 cells as described above. The six CfEcR truncation receptor combinations were duplicated into two groups and treated with either steroid (odd numbers on x-axis of Figure 15) or non-steroid (even numbers on x-axis of Figure 15). In particular, the cells were grown in media containing 0, 1, 5 or 25 uM PonA (steroid) or N-(2-ethyl-3-methoxybenzoyl)-N'-(3,5-dimethylbenzoyl)-N'-tert-butylhydrazine (non-steroid) ligand. The reporter gene activity was measured and total RLU are shown. The number on top of each bar is the maximum fold induction for that treatment and is the mean of three replicates.

As shown in Figure 15, the CfEcRCDEF/MmRXREF receptor combinations were the best switch pairs both in terms of total RLU and fold induction (compare columns 1-6 to columns 7-12). This confirms Applicants' earlier findings as described in Example 2 (Figures 11-14). The same gene expression cassettes encoding the truncated EcR and RXR polypeptides were also assayed in a human lung carcinoma cell line A549 (ATCC) and similar results were observed (data not shown).

25

WE CLAIM:

1. A gene expression modulation system comprising:
  - a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
    - 5 i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated;
    - i) a ligand binding domain comprising a ligand binding domain from a nuclear receptor;
  - b) a second gene expression cassette that is capable of being expressed in  
10 the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
    - i) a transactivation domain; and
    - ii) a ligand binding domain comprising a ligand binding domain from a nuclear receptor other than ultraspiracle (USP);
- 15 wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor, a retinoid X receptor, or an ultraspiracle receptor; and wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.
2. The gene expression modulation system according to claim 1, further comprising a third gene expression cassette comprising:
  - 20 i) a response element to which the DNA-binding domain of the first polypeptide binds;
  - ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
  - iii) the gene whose expression is to be modulated.
- 25 3. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the first polypeptide is an ecdysone receptor polypeptide.
4. The gene expression modulation system according to claim 1, wherein the ligand binding domain of the second polypeptide is a retinoid X receptor polypeptide.
5. A gene expression modulation system comprising:
  - 30 a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:
    - i) a DNA-binding domain that recognizes a response element associated with a gene whose expression is to be modulated; and

- ii) a ligand binding domain comprising a ligand binding domain from an ecdysone receptor; and
- b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:
  - i) a transactivation domain; and
  - ii) a ligand binding domain comprising a ligand binding domain from a retinoid X receptor;

wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

6. The gene expression modulation system according to claim 5, further comprising a third gene expression cassette comprising:

- i) a response element to which the DNA-binding domain of the first polypeptide binds;
- ii) a promoter that is activated by the transactivation domain of the second polypeptide; and
- iii) the gene whose expression is to be modulated.

7. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

8. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

9. The gene expression modulation system according to claim 5, wherein the ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

10. The gene expression modulation system according to claim 5, wherein the

ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

5           11.     A gene expression modulation system comprising:

            a) a first gene expression cassette that is capable of being expressed in a host cell comprising a polynucleotide sequence that encodes a first polypeptide comprising:

- i)    a DNA-binding domain that recognizes a response element  
                      associated with a gene whose expression is to be modulated; and  
10               ii)   a ligand binding domain comprising a ligand binding domain from a  
                      retinoid X receptor; and

            b) a second gene expression cassette that is capable of being expressed in the host cell comprising a polynucleotide sequence that encodes a second polypeptide comprising:

- 15               i)    a transactivation domain; and  
                ii)   a ligand binding domain comprising a ligand binding domain from an  
                      ecdysone receptor;

wherein the ligand binding domains from the first polypeptide and the second polypeptide are different and dimerize.

20           12.     The gene expression modulation system according to claim 11, further comprising a third gene expression cassette comprising:

- i) a response element to which the DNA-binding domain of the first polypeptide binds;  
                ii) a promoter that is activated by the transactivation domain of the second  
25               polypeptide; and  
                iii) the gene whose expression is to be modulated.

            13.     The gene expression modulation system according to claim 11, wherein the ligand binding domain of the first polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22,  
30   SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

            14.     The gene expression modulation system according to claim 11, wherein the ligand binding domain of the first polypeptide comprises an amino acid sequence selected from

the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

15. The gene expression modulation system according to claim 11, wherein the  
5 ligand binding domain of the second polypeptide is encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

16. The gene expression modulation system according to claim 11, wherein the  
10 ligand binding domain of the second polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

17. A gene expression cassette comprising a polynucleotide encoding a hybrid  
15 polypeptide comprising a DNA-binding domain and an ecdysone receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than an ecdysone receptor.

18. The gene expression cassette according to claim 18, wherein the DNA-binding domain is a GAL4 DNA-binding domain or a LexA DNA-binding domain.

20 19. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain and a retinoid X receptor ligand binding domain, wherein the DNA binding domain is from a nuclear receptor other than a retinoid X receptor.

20. The gene expression cassette according to claim 19, wherein the DNA-binding  
25 domain is a GAL4 DNA-binding domain or a LexA DNA-binding domain.

21. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain and an ecdysone receptor ligand binding domain, wherein the transactivation domain is from a nuclear receptor other than an ecdysone receptor.

30 22. The gene expression cassette according to claim 21, wherein the transactivation domain is a VP16 transactivation domain.

23. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain and a retinoid X receptor ligand binding

domain, wherein the transactivation domain is from a nuclear receptor other than a retinoid X receptor.

24. The gene expression cassette according to claim 22, wherein the transactivation domain is a VP16 transactivation domain.

5 25. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ  
10 ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

26. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO:  
15 44) and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

27. A gene expression cassette comprising a polynucleotide encoding a hybrid  
20 polypeptide comprising a DNA-binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 41) or a LexA DBD (SEQ ID NO: 43) and a retinoid X receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID  
25 NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

28. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a DNA-binding domain comprising an amino acid sequence selected from the group consisting of a GAL4 DBD (SEQ ID NO: 42) or a LexA DBD (SEQ ID NO: 44) and a retinoid X receptor ligand binding domain comprising an amino acid sequence  
30 selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

29. A gene expression cassette comprising a polynucleotide encoding a hybrid

polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and an ecdysone receptor ligand binding domain encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, 5 SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

30. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and an ecdysone receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 10 13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, and SEQ ID NO: 20.

31. A gene expression cassette comprising a polynucleotide encoding a hybrid polypeptide comprising a transactivation domain encoded by a polynucleotide comprising a nucleic acid sequence of SEQ ID NO: 45 and a retinoid X receptor ligand binding domain 15 encoded by a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

32. A gene expression cassette comprising a polynucleotide encoding a hybrid 20 polypeptide comprising a transactivation domain comprising an amino acid sequence of SEQ ID NO: 46 and a retinoid X receptor ligand binding domain comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

25 33. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces ligand binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

30 34. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

35. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a

retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation reduces non-steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

36. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
5 retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation mutation enhances ligand binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

37. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation  
10 mutation enhances steroid binding activity of the ecdysone receptor polypeptide or the retinoid X receptor polypeptide.

38. An isolated polynucleotide encoding an ecdysone receptor polypeptide or a  
retinoid X receptor polypeptide comprising a truncation mutation, wherein the truncation  
mutation enhances non-steroid binding activity of the ecdysone receptor polypeptide or the  
15 retinoid X receptor polypeptide.

39. An isolated polynucleotide encoding a retinoid X receptor polypeptide  
comprising a truncation mutation, wherein the truncation mutation increases ligand sensitivity  
of the retinoid X receptor polypeptide.

40. An isolated polynucleotide encoding a retinoid X receptor polypeptide  
20 comprising a truncation mutation, wherein the truncation mutation increases ligand sensitivity  
of a heterodimer, wherein the heterodimer comprises said retinoid X receptor polypeptide and a  
dimerization partner.

41. The isolated polynucleotide according to claim 40, wherein the dimerization  
partner is an ecdysone receptor polypeptide.

25 42. An isolated polynucleotide encoding a truncated ecdysone receptor  
polypeptide, wherein the polynucleotide comprises a nucleic acid sequence selected from the  
group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID  
NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.

43. An isolated polypeptide encoded by the isolated polynucleotide according to  
30 claim 42.

44. An isolated truncated ecdysone receptor polypeptide comprising an amino acid  
sequence selected from the group consisting of SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO:  
13, SEQ ID NO: 14, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18,



SEQ ID NO: 19, and SEQ ID NO: 20.

45. An isolated polynucleotide encoding a truncated retinoid X receptor polypeptide, wherein the polynucleotide comprises a nucleic acid sequence selected from the group consisting of SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, and SEQ ID NO: 30.

46. An isolated polypeptide encoded by the isolated polynucleotide according to claim 45.

47. An isolated truncated retinoid X receptor polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 31, SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 35, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 38, SEQ ID NO: 39, and SEQ ID NO: 40.

48. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

15 a) introducing into the host cell the gene expression modulation system according to claim 1; and


b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

wherein the gene to be expressed is a component of a chimeric gene comprising:

20 i) a response element to which the DNA binding domain from the first polypeptide binds;  
ii) a promoter that is activated by the transactivation domain of the second polypeptide; and  
iii) a gene whose expression is to be modulated,

25 whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

49. The method according to claim 48, wherein the ligand is a compound of the formula:



wherein:

E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

5 R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-  
10 Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

15 R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

20 50. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

a) introducing into the host cell the gene expression modulation system of claim 5; and

25 b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

wherein the gene to be expressed is a component of a chimeric gene comprising:


i) a response element to which the DNA binding domain from the first polypeptide binds;

ii) a promoter that is activated by the transactivation domain of the second polypeptide; and

iii) a gene whose expression is to be modulated,

whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

51. The method according to claim 50, wherein the ligand is a compound of the formula:



wherein:

10 E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

15 R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SEt, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

R<sup>3</sup> is H, Et, or joined with R<sup>2</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

25 R<sup>4</sup>, R<sup>5</sup>, and R<sup>6</sup> are independently H, Me, Et, F, Cl, Br, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CN, C°CH, 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

52. A method of modulating the expression of a gene in a host cell comprising the gene to be modulated comprising the steps of:

a) introducing into the host cell the gene expression modulation system of claim 11; and

b) introducing into the host cell a ligand that independently combines with the ligand binding domains of the first polypeptide and the second polypeptide;

5 wherein the gene to be expressed is a component of a chimeric gene comprising:


i) a response element to which the DNA binding domain from the first polypeptide binds;

ii) a promoter that is activated by the transactivation domain of the second polypeptide; and

10 iii) a gene whose expression is to be modulated,

whereby a complex is formed comprising the ligand, the first polypeptide, and the second polypeptide, and whereby the complex modulates expression of the gene in the host cell.

53. The method according to claim 52, wherein the ligand is a compound of the formula:



15 wherein:

E is a (C<sub>4</sub>-C<sub>6</sub>)alkyl containing a tertiary carbon or a cyano(C<sub>3</sub>-C<sub>5</sub>)alkyl containing a tertiary carbon;

20 R<sup>1</sup> is H, Me, Et, i-Pr, F, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, OH, OMe, OEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, SCN, or SCHF<sub>2</sub>;

25 R<sup>2</sup> is H, Me, Et, n-Pr, i-Pr, formyl, CF<sub>3</sub>, CHF<sub>2</sub>, CHCl<sub>2</sub>, CH<sub>2</sub>F, CH<sub>2</sub>Cl, CH<sub>2</sub>OH, CH<sub>2</sub>OMe, CH<sub>2</sub>CN, CN, C<sup>o</sup>CH, 1-propynyl, 2-propynyl, vinyl, Ac, F, Cl, OH, OMe, OEt, O-n-Pr, OAc, NMe<sub>2</sub>, NEt<sub>2</sub>, SMe, SET, SOCF<sub>3</sub>, OCF<sub>2</sub>CF<sub>2</sub>H, COEt, cyclopropyl, CF<sub>2</sub>CF<sub>3</sub>, CH=CHCN, allyl, azido, OCF<sub>3</sub>, OCHF<sub>2</sub>, O-i-Pr, SCN, SCHF<sub>2</sub>, SMe, NH-CN, or joined with R<sup>3</sup> and the phenyl carbons to which R<sup>2</sup> and R<sup>3</sup> are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

$R^3$  is H, Et, or joined with  $R^2$  and the phenyl carbons to which  $R^2$  and  $R^3$  are attached to form an ethylenedioxy, a dihydrofuryl ring with the oxygen adjacent to a phenyl carbon, or a dihydropyryl ring with the oxygen adjacent to a phenyl carbon;

$R^4$ ,  $R^5$ , and  $R^6$  are independently H, Me, Et, F, Cl, Br, formyl,  $CF_3$ ,  $CHF_2$ ,  $CHCl_2$ ,  
5  $CH_2F$ ,  $CH_2Cl$ ,  $CH_2OH$ , CN,  $C^{\circ}CH$ , 1-propynyl, 2-propynyl, vinyl, OMe, OEt, SMe, or SEt.

54. An isolated host cell into which the gene expression modulation system according to claim 1 has been introduced.

55. The isolated host cell according to claim 54, wherein the host cell is selected  
10 from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell.

56. The isolated host cell according to claim 55, wherein the host cell is a plant cell, a murine cell, or a human cell.

57. An isolated host cell into which the gene expression modulation system  
15 according to claim 5 has been introduced.

58. The isolated host cell according to claim 57, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal cell, and a mammalian cell.

59. The isolated host cell according to claim 58, wherein the host cell is a plant  
20 cell, a murine cell, or a human cell.

60. An isolated host cell into which the gene expression modulation system according to claim 11 has been introduced.

61. The isolated host cell according to claim 60, wherein the host cell is selected from the group consisting of a bacterial cell, a fungal cell, a yeast cell, a plant cell, an animal  
25 cell, and a mammalian cell.

62. The isolated host cell according to claim 61, wherein the host cell is a plant cell, a murine cell, or a human cell.

63. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 1 has been introduced.

30 64. The non-human organism according to claim 63, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal.

65. The non-human organism according to claim 64, wherein the non-human

organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

66. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 5 has been introduced.

5 67. The non-human organism according to claim 66, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an animal, and a mammal.

68. The non-human organism according to claim 67, wherein the non-human organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog,  
10 a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.

69. A non-human organism comprising a host cell into which the gene expression modulation system according to claim 11 has been introduced.

70. The non-human organism according to claim 69, wherein the non-human organism is selected from the group consisting of a bacterium, a fungus, a yeast, a plant, an  
15 animal, and a mammal.

71. The non-human organism according to claim 70, wherein the non-human organism is selected from the group consisting of a plant, a mouse, a rat, a rabbit, a cat, a dog, a bovine, a goat, a pig, a horse, a sheep, a monkey, and a chimpanzee.




Figure 1




Figure 2




Figure 3




Figure 4

**GAL4USP** 

**VP16CfEcR** 

**pGAL4RELuc** 

Figure 5

**GAL4CfEcRVVP16** 

**pGAL4RELuc** 

Figure 6

**VP16CfEcR** 

**pEcRERELuc**  **Luciferase**

Figure 7

**VP16DmEcR** 

**RXR** 


**pE/GRELacZ**  **LacZ**

Figure 8

**VP16CfEcR** 

**RXR** 

**pE/GRELacZ**  **LacZ**

Figure 9

**VP16CfEcR** 

**pE/GRELacZ**  **LacZ**

Figure 10



Analysis of CfEcR Truncations with MmRXRDE in 3T3 Cells




Figure 11

Analysis of CfEcR Truncations with MmRXRE in 3T3 Cells




Figure 12

## Analysis of MmRXR Truncations with CfEcRCDEF in 3T3 Cells




Figure 13

## Analysis of MmRXR Truncations with CfEcRDEF in 3T3 Cells




Figure 14




Figure 15

RH0020.ST25

## SEQUENCE LISTING

<110> Rohm and Haas Company  
 Palli, Subba Reddy  
 Kapitskaya, Marianna Zinovjevna  
 Cress, Dean Ervin

<120> Novel Ecdysone Receptor-Based Inducible Gene Expression System

<130> RH0020

<150> 60/191,355  
 <151> 2000-03-22

<150> 60/269,799  
 <151> 2001-02-20

<160> 64

<170> PatentIn version 3.0

<210> 1  
 <211> 1288  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 1  
 aagggccctg cgccccgtca gcaagaggaa ctgtgtctgg tatgcgggga cagagcctcc 60  
 ggataccact acaatgcgct cacgtgtgaa ggggtgtaaag ggttcttcag acggagtgtt 120  
 accaaaaatg cgggtttatat ttgtaaattc ggtcacgctt gcgaaatgga catgtacatg 180  
 cgacggaaat gccaggagtg ccgcctgaag aagtgccttag ctgtaggcat gaggcctgag 240  
 tgcgtagtac ccgagactca gtgcgccatg aagcggaag agaagaaagc acagaaggag 300  
 aaggacaaac tgctgtctag cacgacgacg gtggacgacc acatgccgcc cattatgcag 360  
 tgtgaacctc cacctcctga agcagcaagg attcacgaag tgggcccaag gtttctctcc 420  
 gacaagctgt tggagacaaa ccggcagaaa aacatcccc agttgacagc caaccagcag 480  
 ttcttatcgc ccaggctcat ctggtaccag gacgggtacg agcagccttc tgatgaagat 540  
 ttgaagagga ttacgcagac gtggcagcaa gcggacgatg aaaacgaaga gtctgacact 600  
 cccttcgcc agatcacaga gatgactatc ctacaggtcc aacttatcgt ggagttcgcg 660  
 aagggattgc cagggttcgc caagatctcg cagcctgac aaattacgct gcttaaggct 720  
 tgctcaagtg aggtaatgat gctccgagtc gcgcgacgat acgatgcggc ctcagacagt 780  
 gttctgttcg cgaacaacca agcgtacact cgcgacaact accgcaaggc tggcatggcc 840  
 tacgtcatcg aggatctact gcacttctgc cgggtgatgt actctatggc gttggacaac 900  
 atccattacg cgctgctcac ggctgtcgtc atcttttctg accggccagg gttggagcag 960  
 ccgcaactgg tggaagaaat ccagcgtac tacctgaata cgctccgcat ctatatcctg 1020

RH0020.ST25

```

aaccagctga gcggttcggc gcgttcgtcc gtcataatacg gcaagatcct ctcaatcctc 1080
tctgagctac gcacgctcgg catgcaaaac tccaacatgt gcatctccct caagctcaag 1140
aacagaaagc tgccgccttt cctcgaggag atctgggatg tggcggacat gtcgcacacc 1200
caaccgccgc ctatcctcga gtccccacg aatctctagc ccctgcgcgc acgcatcgcc 1260
gatgccgcgt ccggccgcgc tgctctga 1288

```

```

<210> 2
<211> 1110
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc feature
<223> Novel Sequence

```

```

<400> 2
gcggtttata tttgtaaatt cggtcacgct tgcgaaatgg acatgtacat gcgacggaaa 60
tgccaggagt gccgcctgaa gaagtgttta gctgtaggca tgaggcctga gtgcgtagta 120
cccgagactc agtgccgcat gaagcggaaa gagaagaaag cacagaagga gaaggacaaa 180
ctgcctgtca gcacgacgac ggtggacgac cacatgccgc ccattatgca gtgtgaacct 240
ccacctctg aagcagcaag gattcacgaa gtggtcccaa ggtttctctc cgacaagctg 300
ttggagacaa accggcagaa aaacatcccc cagttgacag ccaaccagca gttccttate 360
gccaggctca tctggtacca ggacgggtac gagcagcctt ctgatgaaga tttgaagagg 420
attacgcaga cgtggcagca agcggacgat gaaaacgaag agtctgacac tcccttcgcg 480
cagatcacag agatgactat cctcacggtc caacttatcg tggagttcgc gaagggattg 540
ccagggttcg ccaagatctc gcagcctgat caaattacgc tgcttaaggc ttgctcaagt 600
gaggtaatga tgctccgagt cgcgcgacga tacgatgcgg cctcagacag tgttctgttc 660
gcgaacaacc aagcgtacac tcgcgacaac taccgcaagg ctggcatggc ctacgtcatc 720
gaggatctac tgcaacttctg ccggtgcatg tactctatgg cgttggacaa catccattac 780
gcgctgtctc cggtgtctgt catcttttct gaccggccag ggttggagca gccgcaactg 840
gtggaagaaa tccagcggta ctacctgaat acgctccgca tctatatacct gaaccagctg 900
agcgggtcgg cgcgttcgtc cgtcatatac ggcaagatcc tctcaatcct ctctgagcta 960
cgcacgctcg gcatgcaaaa ctccaacatg tgcatctccc tcaagctcaa gaacagaaag 1020
ctgccgcctt tctcagagga gatctgggat gtggcggaca tgtcgcacac ccaaccgccg 1080
cctatcctcg agtccccac gaatctctag 1110

```

```

<210> 3
<211> 1054
<212> DNA

```

RH0020.ST25

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc feature

&lt;223&gt; Novel Sequence

&lt;400&gt; 3

cctgagtgcg tagtaccCGa gactcagtgc gccatgaagc ggaaagagaa gaaagcacag	60
aaggagaagg acaaactgcc tgtcagcacg acgacggtgg acgaccacat gccgcccatt	120
atgcagtgtg aacctccacc tcctgaagca gcaaggattc acgaagtggc cccaaggttt	180
ctctccgaca agctgttgga gacaaaccgg cagaaaaaca tccccagtt gacagccaac	240
cagcagttcc ttatcgccag gctcatctgg taccaggacg ggtacgagca gccttctgat	300
gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct	360
gacactccct tccgccagat cacagagatg actatcctca cgggccaaact tatcgtggag	420
ttcgCGaagg gattgccagg gttcgccaag atctcgCagc ctgatcaaatacgtctgctt	480
aaggcttgct caagtGagggt aatgatgctc cgagtcgcgc gacgatacga tgcggcctca	540
gacagtgttc tgttcgcgaa caaccaagcg tacactcgcg acaactaccg caaggctggc	600
atggcctacg tcatcgagga tctactgcac ttctgccggg gcatgtactc tatggcgcttg	660
gacaacatcc attacgcgct gctcacggct gtcgtcatct tttctgaccg gccagggttg	720
gagcagccgc aactggtgga agaaatccag cggctactacc tgaatacgct ccgcatttat	780
atcctgaacc agctgagcgg gtcggcgcggt tcgtccgtca tatacggcaa gatcctctca	840
atcctctctg agctacgcac gtcggcatg caaaactcca acatgtgcat ctccctcaag	900
ctcaagaaca gaaagctgcc gcctttcctc gaggagatct gggatgtggc ggacatgtcg	960
cacacccaac cgcgccttat cctcgagtc cccacgaatc tctagcccct gcgcgcacgc	1020
atcgccgatg ccgcgtccgg ccgcgtgct ctga	1054

&lt;210&gt; 4

&lt;211&gt; 735

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc feature

&lt;223&gt; Novel Sequence

&lt;400&gt; 4

taccaggacg ggtacgagca gccttctgat gaagatttga agaggattac gcagacgtgg	60
cagcaagcgg acgatgaaaa cgaagagtct gacactccct tccgccagat cacagagatg	120
actatcctca cgggccaaact tatcgtggag ttcgCGaagg gattgccagg gttcgccaag	180
atctcgCagc ctgatcaaatacgtctgctt aaggcttgct caagtGagggt aatgatgctc	240
cgagtcgcgc gacgatacga tgcggcctca gacagtgttc tgttcgcgaa caaccaagcg	300

RH0020.ST25

```

tacactcgcg acaactaccg caaggctggc atggcctacg tcatcgagga tctactgcac 360
ttctgcccgt gcatgtactc tatggcggtg gacaacatcc attacgcgct gtcacggct 420
gtcgtcatct tttctgaccg gccagggttg gagcagccgc aactggtgga agaaatccag 480
cggtactacc tgaatacgct ccgcatctat atcctgaacc agctgagcgg gtcggcgcg 540
tcgtccgtca tatacggcaa gatcctctca atcctctctg agctacgcac gtcggcatg 600
caaaactcca acatgtgcat ctccctcaag ctcaagaaca gaaagctgcc gcctttctc 660
gaggagatct gggatgtggc ggacatgtcg cacaccaac cgccgcctat cctcgagtcc 720
cccaogaatc tctag 735

```

```

<210> 5
<211> 960
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc_feature
<223> Novel Sequence

```

```

<400> 5
cctgagtgcg tagtaccoga gactcagtgc gccatgaagc ggaaagagaa gaaagcacag 60
aaggagaagg aaaaactgcc tgtcagcacg acgacgggtg acgaccacat gccgcccatt 120
atgcagtgtg aacctccacc tcctgaagca gcaaggattc acgaagtggc cccaaggttt 180
ctctccgaca agctgttgga gacaaaccgg cagaaaaaca tccccagtt gacagccaac 240
cagcagttcc ttatcgccag gctcatctgg taccaggacg ggtacgagca gccttctgat 300
gaagatttga agaggattac gcagacgtgg cagcaagcgg acgatgaaaa cgaagagtct 360
gacactccct tccgccagat cacagagatg actatcctca cggccaact tatcgtggag 420
ttcgcgaagg gattgccagg gttcgccaag atctcgagc ctgatcaaat tacgctgctt 480
aaggcttgct caagtgaggt aatgatgctc cgagtgcgcg gacgatacga tgcggcctca 540
gacagtgttc tgttcgcgaa caaccaagcg tacactcgcg acaactaccg caaggctggc 600
atggcctacg tcatcgagga tctactgcac ttctgcccgt gcatgtactc tatggcggtg 660
gacaacatcc attacgcgct gtcacggct gtcgtcatct tttctgaccg gccagggttg 720
gagcagccgc aactggtgga agaaatccag cgggtactacc tgaatacgct ccgcatctat 780
atcctgaacc agctgagcgg gtcggcgcg  tcgtccgtca tatacggcaa gatcctctca 840
atcctctctg agctacgcac gtcggcatg caaaactcca acatgtgcat ctccctcaag 900
ctcaagaaca gaaagctgcc gcctttctc gaggagatct gggatgtggc ggacatgtcg 960

```

```

<210> 6
<211> 1878
<212> DNA

```

RH0020.ST25

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; misc feature

&lt;223&gt; Novel Sequence

&lt;400&gt; 6

```

ggacctgcgc cacgggtgca agaggagctg tgcctggttt gcggcgacag ggacctccggc      60
taccactaca acgccctcac ctgtgagggc tgcaaggggt tctttcgacg cagcgttacg      120
aagagcgccg tctactgctg caagttcggg cgcgcctgcg aaatggacat gtacatgagg      180
cgaaagtgtc aggagtgccg cctgaaaaag tgcctggccg tgggtatgcg gccggaatgc      240
gtcgtcccg agaaccaatg tgcgatgaag cggcgcgaaa agaaggcca gaaggagaag      300
gacaaaatga ccacttcgcc gagctctcag catggcggca atggcagctt ggctcttgt      360
ggcggccaaag actttgttaa gaaggagatt cttgacctta tgacatgcga gccgcccag      420
catgccacta ttccgctact acctgatgaa atattggcca agtgtcaagc gcgcaatata      480
ccttccttaa cgtacaatca gttggcgtt atatacaagt taatttggtta ccaggatggc      540
tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccga tgagaacgag      600
agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg      660
attgttgagt ttgctaaagg tctaccagcg tttaaaaga taccacagga ggaccagatc      720
acgttactaa aggcctgctc gtcggaggtg atgatgtgc gtatggcacg acgtatgac      780
cacagctcgg actcaatatt cttcggaat aatagatcat atacgcgga ttcttcaaaa      840
atggccggaa tggctgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg      900
atgaaggtgg acaacgtcga atacgcgctt ctactgcca ttgtgatctt ctcggaccgg      960
ccgggcctgg agaaggcca actagtgcga gcgatccaga gctactacat cgacacgcta     1020
cgcatttata tactcaaccg ccactgcggc gactcaatga gcctcgtctt ctacgcaaag     1080
ctgctctcga tctcaccga gctgcgtacg ctgggcaacc agaacgccga gatgtgtttc     1140
tcactaaagc tcaaaaaccg caaactgcc aagttcctcg aggagatctg ggacgttcat     1200
gccatcccg catcggtcca gtcgcacctt cagattacc aggaggagaa cgagcgtctc     1260
gagcgggctg agcgtatgcg gccatcgggtt gggggcgcca ttaccgccg cattgattgc     1320
gactctgcct ccacttcggc ggcggcagcc gcggcccagc atcagcctca gcctcagccc     1380
cagccccaac cctcctccct gaccagaaac gattcccagc accagacaca gccgcagcta     1440
caacctcagc taccacctca gctgcaaggc caactgcaac ccagctcca accacagctt     1500
cagacgcaac tccagccaca gattcaacca cagccacagc tccttcccg ctccgctccc     1560
gtgcccgcct ccgtaaccgc acctggttcc ttgtccgcgg tcagtacgag cagcgaatac     1620
atgggcggaa gtgcggccat aggaccatc acgccggcaa ccaccagcag tatcacggct     1680
gccgttacgg ctagctccac cacatcagcg gtaccgatgg gcaacggagt tggagtcgg      1740

```



## RH0020.ST25

gttgggggtgg gcggaacgt cagcatgtat gcgaacgccc agacggcgat ggccttgatg 1800  
 ggtgtagccc tgcattcgca ccaagagcag cttatcgggg gagtggcggg taagtcggag 1860  
 cactcgacga ctgcatag 1878

<210> 7  
 <211> 1752  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 7  
 gccgtctact gctgcaagtt cgggcgcgcc tgcgaaatgg acatgtacat gaggcgaaag 60  
 tgtcaggagt gccgcctgaa aaagtgcctg gccgtgggta tgcggccgga atgcgtcgtc 120  
 ccggagaacc aatgtgcgat gaagcggcgc gaaaagaagg ccagaagga gaaggacaaa 180  
 atgaccactt cgccgagctc tcagcatggc ggcaatggca gcttggcctc tgggtggcggc 240  
 caagactttg ttaagaagga gattcttgac cttatgacat gcgagccgcc ccagcatgcc 300  
 actattccgc tactacctga tgaaatattg gccaaagtgc aagcgcgcaa tataccttcc 360  
 ttaacgtaca atcagttggc cgttatatac aagttaattt ggtaccagga tggctatgag 420  
 cagccatctg aagaggatct caggcgtata atgagtcaac ccgatgagaa cgagagccaa 480  
 acggacgtca gctttcggca tataaccgag ataaccatac tcacgggtcca gttgattgtt 540  
 gagtttgcta aaggtctacc agcgtttaca aagatacccc aggaggacca gatcacgtta 600  
 ctaaaggcct gctcgtcggga ggtgatgatg ctgcgtatgg cagcagccta tgaccacagc 660  
 tcggactcaa tattcttcgc gaataataga tcatatacgc gggattctta caaatggcc 720  
 ggaatggctg ataacattga agacctgctg cttttctgcc gccaaatgtt ctcgatgaag 780  
 gtggacaacg tcgaatacgc gcttctcact gccattgtga ttttctcgga ccggccgggc 840  
 ctggagaagg cccaactagt cgaagcgatc cagagctact acatcgacac gctacgcatt 900  
 tatatactca accgccactg cggcgactca atgagcctcg ttttctacgc aaagctgctc 960  
 tcgactctca ccgagctgcg tacgctgggc aaccagaacg ccgagatgtg tttctcacta 1020  
 aagctcaaaa accgcaaact gcccaagttc ctcgaggaga tctgggacgt tcatgccatc 1080  
 ccgccatcgg tccagtcgca ccttcagatt acccaggagg agaacgagcg tctcgagcgg 1140  
 gctgagcgtg tgcgggcacg ggttgggggc gccattaccg ccggcattga ttgcgactct 1200  
 gcctccactt cggcggcggc agccgcggcc cagcatcagc ctcagcctca gcccagccc 1260  
 caaccctcct ccctgaccca gaacgattcc cagcaccaga cacagccgca gctacaacct 1320  
 cagctaccac ctcagctgca aggtcaactg caacccagc tccaaccaca gcttcagacg 1380

RH0020.ST25

caactccagc cacagattca accacagcca cagctccttc ccgtctccgc tcccgtgccc	1440
gcctccgtaa ccgcacctgg ttctttgtcc gcggtcagta cgagcagcga atacatgggc	1500
ggaagtgcgg ccataggacc catcacgccg gcaaccacca gcagtatcac ggctgccgtt	1560
accgctagct ccaccacatc agcgggtaccg atgggcaacg gagttggagt cgggtgttggg	1620
gtgggcgcca acgtcagcat gtatgcgaac gccagacgg cgatggcctt gatgggtgta	1680
gccctgcatt cgcaccaaga gcagcttatc gggggagtgg cggttaagtc ggagcactcg	1740
acgactgcat ag	1752

<210> 8  
 <211> 1650  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 8	
cggccggaat gcgtcgtccc ggagaaccaa tgtgcgatga agcggcgcgga aaagaaggcc	60
cagaaggaga aggacaaaat gaccacttcg ccgagctctc agcatggcgg caatggcagc	120
ttggcctctg gtggcggcca agactttgtt aagaaggaga ttcttgacct tatgacatgc	180
gagccgcccc agcatgccac tattccgcta ctacctgatg aaatattggc caagtgtcaa	240
gcgcgcaata taccttcctt aacgtacaat cagttggccg ttatatacaa gttaatttgg	300
taccaggatg gctatgagca gccatctgaa gaggatctca ggcgtataat gagtcaaccc	360
gatgagaacg agagccaaac ggacgtcagc tttcggcata taaccgagat aaccatactc	420
acggtccagt tgattgttga gtttgctaaa ggtctaccag cgtttacaaa gataccccag	480
gaggaccaga tcacgttact aaaggcctgc tcgtcggagg tgatgatgct gcgtatggca	540
cgacgctatg accacagctc ggactcaata ttcttcgcca ataatagatc atatacggcg	600
gattcttaca aaatggccgg aatggctgat aacattgaag acctgctgca tttctgccgc	660
caaatgttct cgatgaagggt ggacaacgtc gaatacgcgc ttctcactgc cattgtgatc	720
ttctcggacc ggccgggcct ggagaaggcc caactagtcg aagcgatcca gagctactac	780
atcgacacgc tacgcattta tatactcaac cgccactgcg gcgactcaat gaggcctcgtc	840
ttctacgcaa agctgctctc gatcctcacc gagctgcgta cgctgggcaa ccagaacgcc	900
gagatgtgtt tctcactaaa gctcaaaaac cgcaaactgc ccaagttcct cgaggagatc	960
tgggacgttc atgcatccc gccatcggtc cagtcgcacc ttcagattac ccaggaggag	1020
aacgagcgtc tcgagcgggc tgagcgtatg cgggcatcgg ttggggggcg cattaccgcc	1080
ggcattgatt gcgactctgc ctccacttcg gcggcggcag ccgcggccca gcatcagcct	1140
cagcctcagc ccagcccca accctcctcc ctgaccacga acgattccca gcaccagaca	1200

## RH0020.ST25

```

cagccgcagc tacaacctca gctaccacct cagctgcaag gtcaactgca accccagctc 1260
caaccacagc ttccagacgca actccagcca cagattcaac cacagccaca gtccttccc 1320
gtctccgctc ccgtgcccgc ctccgtaacc gcacctggtt ccttgctccg ggtcagtagc 1380
agcagcgaat acatgggagg aagtgcgggc ataggaccca tcacgccggc aaccaccagc 1440
agtatcacgg ctgccgttac cgctagctcc accacatcag cggtagcgat gggcaacgga 1500
gttgaggatc gtgttggggt gggcggaac gtcagcatgt atgcgaacgc ccagacggcg 1560
atggccttga tgggtgtagc cctgcattcg caccaagagc agcttatcgg gggagtggcg 1620
gttaagtcgg agcactcgac gactgcatag 1650

```

```

<210> 9
<211> 1338
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc_feature
<223> Novel Sequence

```

```

<400> 9
tatgagcagc catctgaaga ggatctcagg cgtataatga gtcaaccgga tgagaacgag 60
agccaaacgg acgtcagctt tcggcatata accgagataa ccatactcac ggtccagttg 120
attgttgagt ttgctaaagg tctaccagcg ttacaaaaga taccacagga ggaccagatc 180
acgttactaa aggcctgctc gtcggagggtg atgatgctgc gtatggcacg acgctatgac 240
cacagctcgg actcaatatt cttcggaat aatagatcat atacgcggga ttcttataaa 300
atggccggaa tggctgataa cattgaagac ctgctgcatt tctgccgcca aatgttctcg 360
atgaaggtag acaacgtcga atacgcgctt ctactgcca ttgtgatctt ctggaccgg 420
ccgggcctgg agaaggccca actagtcgaa gcgatccaga gctactacat cgacacgcta 480
cgcatttata tactcaaccg cactgcgggc gactcaatga gctcgtctt ctacgaaaag 540
ctgctctcga tcctcacgga gctgcgtag ctgggcaacc agaacgccga gatgtgtttc 600
tcactaaagc taaaaaacg caaactgcc aagttcctcg aggagatctg ggacgttcat 660
gccatcccgc catcgggtcca gtcgcacct cagattaccc aggaggagaa cgagcgtctc 720
gagcgggctg agcgtatgag ggcacgggtt gggggcgcca ttaccgccgg cattgattgc 780
gactctgcct ccacttcggc ggcggcagcc gcggccagc atcagcctca gcctcagccc 840
cagccccaac cctcctcctt gaccagaaac gattcccagc accagacaca gccgcagcta 900
caacctcagc taccacctca gctgcaaggt caactgcaac ccagctcca accacagctt 960
cagacgcaac tccagccaca gattcaacca cagccacagc tccttcccgt ctccgctccc 1020
gtgccgcct ccgtaaccgc acctgggttc ttgtccgcgg tcagtacgag cagcgaatac 1080

```

RH0020.ST25

atgggcgga gtcgggcat aggacccatc acgccggcaa ccaccagcag tatcacggct 1140  
gccgttaccg ctagctccac cacatcagcg gtaccgatgg gcaacggagt tggagtcggg 1200  
gttggggtgg gcggcaacgt cagcatgtat gcgaacgcc agacggcgat ggccttgatg 1260  
ggtgtagccc tgcattcgca ccaagagcag cttatcgggg gagtggcggg taagtcggag 1320  
cactcgacga ctgcatag 1338

<210> 10  
<211> 969  
<212> DNA  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

<400> 10  
cggccggaat gcgtcgtccc ggagaaccaa tgtgcgatga agcggcgcgga aaagaaggcc 60  
cagaaggaga aggacaaaat gaccacttcg ccgagctctc agcatggcgg caatggcagc 120  
ttggcctctg gtggcggcca agactttgtt aagaaggaga ttcttgacct tatgacatgc 180  
gagccgcccc agcatgccac tattccgcta ctacctgatg aaatattggc caagtgtcaa 240  
gcgcgcaata taccttcctt aacgtacaat cagttggccg ttatatacaa gttaatttgg 300  
taccaggatg gctatgagca gccatctgaa gaggatctca ggcgtataat gagtcaaccc 360  
gatgagaacg agagccaaac ggacgtcagc ttctcgcata taaccgagat aaccatactc 420  
acggtccagt tgattgttga gtttgctaaa ggtctaccag cgtttacaaa gataccccag 480  
gaggaccaga tcacgttact aaaggcctgc tcgtcggagg tgatgatgct gcgtatggca 540  
cgacgctatg accacagctc ggactcaata ttcttcgcga ataatagatc atatacgcg 600  
gattcttaca aaatggccgg aatggctgat aacattgaag acctgctgca tttctgccgc 660  
caaatgttct cgatgaaggt ggacaacgtc gaatacgcgc ttctcactgc cattgtgatc 720  
ttctcggacc ggccgggcct ggagaaggcc caactagtcg aagcgatcca gagctactac 780  
atcgacacgc tacgcattta tatactcaac cgccactgcg gcgactcaat gagcctcgtc 840  
ttctacgcaa agctgctctc gatcctcacc gagctgcgta cgctgggcaa ccagaacgcc 900  
gagatgtgtt tctcactaaa gctcaaaaac cgaaaactgc ccaagttcct cgaggagatc 960  
tgggacgtt 969

<210> 11  
<211> 412  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

RH0020.ST25

&lt;400&gt; 11

Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly  
 1 5 10 15  
 Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys  
 20 25 30  
 Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys  
 35 40 45  
 Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys  
 50 55 60  
 Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu  
 65 70 75 80  
 Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys  
 85 90 95  
 Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp  
 100 105 110  
 Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro Glu Ala  
 115 120 125  
 Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu  
 130 135 140  
 Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln  
 145 150 155 160  
 Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro  
 165 170 175  
 Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp  
 180 185 190  
 Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met  
 195 200 205  
 Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro  
 210 215 220  
 Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala  
 225 230 235 240  
 Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala  
 245 250 255  
 Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp  
 260 265 270  
 Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His  
 275 280 285  
 Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala  
 290 295 300  
 Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln  
 305 310 315 320  
 Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg

RH0020.ST25

```

          325                      330                      335
Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile
          340                      345                      350
Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met
          355                      360                      365
Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu
          370                      375                      380
Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr
          385                      390                      395                      400
Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu
          405                      410

```

```

<210> 12
<211> 412
<212> PRT
<213> Artificial Sequence

```

```

<220>
<221> misc feature
<223> Novel Sequence

```

```

<400> 12

```

```

Lys Gly Pro Ala Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly
1          5          10          15
Asp Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys
          20          25          30
Lys Gly Phe Phe Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys
          35          40          45
Lys Phe Gly His Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys
          50          55          60
Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu
          65          70          75          80
Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys
          85          90          95
Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp
          100          105          110
Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro Glu Ala
          115          120          125
Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu
          130          135          140
Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln
          145          150          155          160
Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro
          165          170          175
Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp
          180          185          190

```

RH0020.ST25

Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met  
 195 200 205  
 Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro  
 210 215 220  
 Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu Lys Ala  
 225 230 235 240  
 Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala  
 245 250 255  
 Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp  
 260 265 270  
 Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His  
 275 280 285  
 Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala  
 290 295 300  
 Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln  
 305 310 315 320  
 Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg  
 325 330 335  
 Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile  
 340 345 350  
 Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met  
 355 360 365  
 Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu  
 370 375 380  
 Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr  
 385 390 395 400  
 Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu  
 405 410

<210> 13  
 <211> 334  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 13

Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu  
 1 5 10 15  
 Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr  
 20 25 30  
 Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro Pro  
 35 40 45  
 Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys  
 50 55 60

RH0020.ST25

Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn  
 65 70 75 80  
 Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu  
 85 90 95  
 Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln  
 100 105 110  
 Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr  
 115 120 125  
 Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly  
 130 135 140  
 Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu  
 145 150 155 160  
 Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr  
 165 170 175  
 Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr  
 180 185 190  
 Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu  
 195 200 205  
 Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His  
 210 215 220  
 Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu  
 225 230 235 240  
 Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr  
 245 250 255  
 Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser  
 260 265 270  
 Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu  
 275 280 285  
 Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg  
 290 295 300  
 Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser  
 305 310 315 320  
 His Thr Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn Leu  
 325 330

<210> 14  
 <211> 244  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 14

Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile



RH0020.ST25

1	5	10	15
Thr Gln Thr Trp Gln Gln Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr	20	25	30
Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile	35	40	45
Val Glu Phe Ala Lys Gly Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro	50	55	60
Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu	65	70	75
Arg Val Ala Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val Leu Phe Ala	85	90	95
Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala	100	105	110
Tyr Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met	115	120	125
Ala Leu Asp Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe	130	135	140
Ser Asp Arg Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln	145	150	155
Arg Tyr Tyr Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser	165	170	175
Gly Ser Ala Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu	180	185	190
Ser Glu Leu Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser	195	200	205
Leu Lys Leu Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp	210	215	220
Asp Val Ala Asp Met Ser His Thr Gln Pro Pro Pro Ile Leu Glu Ser	225	230	235
Pro Thr Asn Leu			

<210> 15  
 <211> 320  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 15

Pro Glu Cys Val Val Pro Glu Thr Gln Cys Ala Met Lys Arg Lys Glu	1	5	10	15
Lys Lys Ala Gln Lys Glu Lys Asp Lys Leu Pro Val Ser Thr Thr Thr	20	25	30	

RH0020.ST25

Val Asp Asp His Met Pro Pro Ile Met Gln Cys Glu Pro Pro Pro  
 35 40 45  
 Glu Ala Ala Arg Ile His Glu Val Val Pro Arg Phe Leu Ser Asp Lys  
 50 55 60  
 Leu Leu Glu Thr Asn Arg Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn  
 65 70 75 80  
 Gln Gln Phe Leu Ile Ala Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu  
 85 90 95  
 Gln Pro Ser Asp Glu Asp Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln  
 100 105 110  
 Ala Asp Asp Glu Asn Glu Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr  
 115 120 125  
 Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly  
 130 135 140  
 Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile Thr Leu Leu  
 145 150 155 160  
 Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala Arg Arg Tyr  
 165 170 175  
 Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr  
 180 185 190  
 Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile Glu Asp Leu  
 195 200 205  
 Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp Asn Ile His  
 210 215 220  
 Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg Pro Gly Leu  
 225 230 235 240  
 Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr  
 245 250 255  
 Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser  
 260 265 270  
 Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu  
 275 280 285  
 Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu Lys Asn Arg  
 290 295 300  
 Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala Asp Met Ser  
 305 310 315 320

<210> 16  
 <211> 625  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 16

RH0020.ST25

Gly Pro Ala Pro Arg Val Gln Glu Glu Leu Cys Leu Val Cys Gly Asp  
 1 5 10 15  
 Arg Ala Ser Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys  
 20 25 30  
 Gly Phe Phe Arg Arg Ser Val Thr Lys Ser Ala Val Tyr Cys Cys Lys  
 35 40 45  
 Phe Gly Arg Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln  
 50 55 60  
 Glu Cys Arg Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys  
 65 70 75 80  
 Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg Glu Lys Lys Ala  
 85 90 95  
 Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser Ser Gln His Gly  
 100 105 110  
 Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp Phe Val Lys Lys  
 115 120 125  
 Glu Ile Leu Asp Leu Met Thr Cys Glu Pro Pro Gln His Ala Thr Ile  
 130 135 140  
 Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys Cys Gln Ala Arg Asn Ile  
 145 150 155 160  
 Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val Ile Tyr Lys Leu Ile Trp  
 165 170 175  
 Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile  
 180 185 190  
 Met Ser Gln Pro Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg  
 195 200 205  
 His Ile Thr Glu Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe  
 210 215 220  
 Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile  
 225 230 235 240  
 Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala  
 245 250 255  
 Arg Arg Tyr Asp His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg  
 260 265 270  
 Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile  
 275 280 285  
 Glu Asp Leu Leu His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp  
 290 295 300  
 Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg  
 305 310 315 320  
 Pro Gly Leu Glu Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr  
 325 330 335  
 Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser

RH0020.ST25

```

      340              345              350
Met Ser Leu Val Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu
      355              360              365
Arg Thr Leu Gly Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu
      370              375              380
Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His
      385              390              395              400
Ala Ile Pro Pro Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu
      405              410              415
Asn Glu Arg Leu Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly
      420              425              430
Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala
      435              440              445
Ala Ala Ala Ala Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro
      450              455              460
Ser Ser Leu Thr Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu
      465              470              475              480
Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu
      485              490              495
Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro
      500              505              510
Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro
      515              520              525
Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser
      530              535              540
Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala
      545              550              555              560
Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly
      565              570              575
Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn
      580              585              590
Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu His Ser His Gln
      595              600              605
Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr
      610              615              620

```

Ala  
625

<210> 17  
<211> 583  
<212> PRT  
<213> Artificial Sequence

<220>  
<221> misc\_feature  
<223> Novel Sequence

RH0020.ST25

&lt;400&gt; 17

Ala Val Tyr Cys Cys Lys Phe Gly Arg Ala Cys Glu Met Asp Met Tyr  
 1 5 10 15  
 Met Arg Arg Lys Cys Gln Glu Cys Arg Leu Lys Lys Cys Leu Ala Val  
 20 25 30  
 Gly Met Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys  
 35 40 45  
 Arg Arg Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser  
 50 55 60  
 Pro Ser Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly  
 65 70 75 80  
 Gln Asp Phe Val Lys Lys Glu Ile Leu Asp Leu Met Thr Cys Glu Pro  
 85 90 95  
 Pro Gln His Ala Thr Ile Pro Leu Leu Pro Asp Glu Ile Leu Ala Lys  
 100 105 110  
 Cys Gln Ala Arg Asn Ile Pro Ser Leu Thr Tyr Asn Gln Leu Ala Val  
 115 120 125  
 Ile Tyr Lys Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Glu  
 130 135 140  
 Glu Asp Leu Arg Arg Ile Met Ser Gln Pro Asp Glu Asn Glu Ser Gln  
 145 150 155 160  
 Thr Asp Val Ser Phe Arg His Ile Thr Glu Ile Thr Ile Leu Thr Val  
 165 170 175  
 Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro Ala Phe Thr Lys Ile  
 180 185 190  
 Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys Ala Cys Ser Ser Glu Val  
 195 200 205  
 Met Met Leu Arg Met Ala Arg Arg Tyr Asp His Ser Ser Asp Ser Ile  
 210 215 220  
 Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg Asp Ser Tyr Lys Met Ala  
 225 230 235 240  
 Gly Met Ala Asp Asn Ile Glu Asp Leu Leu His Phe Cys Arg Gln Met  
 245 250 255  
 Phe Ser Met Lys Val Asp Asn Val Glu Tyr Ala Leu Leu Thr Ala Ile  
 260 265 270  
 Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Lys Ala Gln Leu Val Glu  
 275 280 285  
 Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu Arg Ile Tyr Ile Leu Asn  
 290 295 300  
 Arg His Cys Gly Asp Ser Met Ser Leu Val Phe Tyr Ala Lys Leu Leu  
 305 310 315 320  
 Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly Asn Gln Asn Ala Glu Met  
 325 330 335

## RH0020.ST25

Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Lys Phe Leu Glu  
 340 345 350  
 Glu Ile Trp Asp Val His Ala Ile Pro Pro Ser Val Gln Ser His Leu  
 355 360 365  
 Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu Glu Arg Ala Glu Arg Met  
 370 375 380  
 Arg Ala Ser Val Gly Gly Ala Ile Thr Ala Gly Ile Asp Cys Asp Ser  
 385 390 395 400  
 Ala Ser Thr Ser Ala Ala Ala Ala Ala Gln His Gln Pro Gln Pro  
 405 410 415  
 Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr Gln Asn Asp Ser Gln His  
 420 425 430  
 Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly  
 435 440 445  
 Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro  
 450 455 460  
 Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro  
 465 470 475 480  
 Ala Ser Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser  
 485 490 495  
 Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr  
 500 505 510  
 Thr Ser Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala  
 515 520 525  
 Val Pro Met Gly Asn Gly Val Gly Val Gly Val Gly Gly Asn  
 530 535 540  
 Val Ser Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val  
 545 550 555 560  
 Ala Leu His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys  
 565 570 575  
 Ser Glu His Ser Thr Thr Ala  
 580

<210> 18  
 <211> 549  
 <212> PRT  
 <213> Artificial Sequence

<400> 18

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg  
 1 5 10 15  
 Glu Lys Lys Ala Gln Lys Glu Lys Asp Lys Met Thr Thr Ser Pro Ser  
 20 25 30  
 Ser Gln His Gly Gly Asn Gly Ser Leu Ala Ser Gly Gly Gly Gln Asp  
 35 40 45

## RH0020.ST25

Phe	Val	Lys	Lys	Glu	Ile	Leu	Asp	Leu	Met	Thr	Cys	Glu	Pro	Pro	Gln
50						55					60				
His	Ala	Thr	Ile	Pro	Leu	Leu	Pro	Asp	Glu	Ile	Leu	Ala	Lys	Cys	Gln
65					70				75						80
Ala	Arg	Asn	Ile	Pro	Ser	Leu	Thr	Tyr	Asn	Gln	Leu	Ala	Val	Ile	Tyr
			85						90					95	
Lys	Leu	Ile	Trp	Tyr	Gln	Asp	Gly	Tyr	Glu	Gln	Pro	Ser	Glu	Glu	Asp
			100					105					110		
Leu	Arg	Arg	Ile	Met	Ser	Gln	Pro	Asp	Glu	Asn	Glu	Ser	Gln	Thr	Asp
		115					120					125			
Val	Ser	Phe	Arg	His	Ile	Thr	Glu	Ile	Thr	Ile	Leu	Thr	Val	Gln	Leu
	130					135					140				
Ile	Val	Glu	Phe	Ala	Lys	Gly	Leu	Pro	Ala	Phe	Thr	Lys	Ile	Pro	Gln
145					150					155					160
Glu	Asp	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Cys	Ser	Ser	Glu	Val	Met	Met
				165					170					175	
Leu	Arg	Met	Ala	Arg	Arg	Tyr	Asp	His	Ser	Ser	Asp	Ser	Ile	Phe	Phe
			180					185					190		
Ala	Asn	Asn	Arg	Ser	Tyr	Thr	Arg	Asp	Ser	Tyr	Lys	Met	Ala	Gly	Met
		195					200					205			
Ala	Asp	Asn	Ile	Glu	Asp	Leu	Leu	His	Phe	Cys	Arg	Gln	Met	Phe	Ser
	210					215					220				
Met	Lys	Val	Asp	Asn	Val	Glu	Tyr	Ala	Leu	Leu	Thr	Ala	Ile	Val	Ile
225					230					235					240
Phe	Ser	Asp	Arg	Pro	Gly	Leu	Glu	Lys	Ala	Gln	Leu	Val	Glu	Ala	Ile
				245					250					255	
Gln	Ser	Tyr	Tyr	Ile	Asp	Thr	Leu	Arg	Ile	Tyr	Ile	Leu	Asn	Arg	His
			260					265					270		
Cys	Gly	Asp	Ser	Met	Ser	Leu	Val	Phe	Tyr	Ala	Lys	Leu	Leu	Ser	Ile
		275					280					285			
Leu	Thr	Glu	Leu	Arg	Thr	Leu	Gly	Asn	Gln	Asn	Ala	Glu	Met	Cys	Phe
	290					295					300				
Ser	Leu	Lys	Leu	Lys	Asn	Arg	Lys	Leu	Pro	Lys	Phe	Leu	Glu	Glu	Ile
305					310					315					320
Trp	Asp	Val	His	Ala	Ile	Pro	Pro	Ser	Val	Gln	Ser	His	Leu	Gln	Ile
				325					330					335	
Thr	Gln	Glu	Glu	Asn	Glu	Arg	Leu	Glu	Arg	Ala	Glu	Arg	Met	Arg	Ala
			340					345					350		
Ser	Val	Gly	Gly	Ala	Ile	Thr	Ala	Gly	Ile	Asp	Cys	Asp	Ser	Ala	Ser
		355					360					365			
Thr	Ser	Ala	Ala	Ala	Ala	Ala	Ala	Gln	His	Gln	Pro	Gln	Pro	Gln	Pro
	370					375					380				
Gln	Pro	Gln	Pro	Ser	Ser	Leu	Thr	Gln	Asn	Asp	Ser	Gln	His	Gln	Thr
385					390					395					400

RH0020.ST25

Gln Pro Gln Leu Gln Pro Gln Leu Pro Pro Gln Leu Gln Gly Gln Leu  
 405 410 415  
 Gln Pro Gln Leu Gln Pro Gln Leu Gln Thr Gln Leu Gln Pro Gln Ile  
 420 425 430  
 Gln Pro Gln Pro Gln Leu Leu Pro Val Ser Ala Pro Val Pro Ala Ser  
 435 440 445  
 Val Thr Ala Pro Gly Ser Leu Ser Ala Val Ser Thr Ser Ser Glu Tyr  
 450 455 460  
 Met Gly Gly Ser Ala Ala Ile Gly Pro Ile Thr Pro Ala Thr Thr Ser  
 465 470 475 480  
 Ser Ile Thr Ala Ala Val Thr Ala Ser Ser Thr Thr Ser Ala Val Pro  
 485 490 495  
 Met Gly Asn Gly Val Gly Val Gly Val Gly Val Gly Gly Asn Val Ser  
 500 505 510  
 Met Tyr Ala Asn Ala Gln Thr Ala Met Ala Leu Met Gly Val Ala Leu  
 515 520 525  
 His Ser His Gln Glu Gln Leu Ile Gly Gly Val Ala Val Lys Ser Glu  
 530 535 540  
 His Ser Thr Thr Ala  
 545

<210> 19  
 <211> 445  
 <212> PRT  
 <213> Artificial Sequence  
 <400> 19

Tyr Glu Gln Pro Ser Glu Glu Asp Leu Arg Arg Ile Met Ser Gln Pro  
 1 5 10 15  
 Asp Glu Asn Glu Ser Gln Thr Asp Val Ser Phe Arg His Ile Thr Glu  
 20 25 30  
 Ile Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu  
 35 40 45  
 Pro Ala Phe Thr Lys Ile Pro Gln Glu Asp Gln Ile Thr Leu Leu Lys  
 50 55 60  
 Ala Cys Ser Ser Glu Val Met Met Leu Arg Met Ala Arg Arg Tyr Asp  
 65 70 75 80  
 His Ser Ser Asp Ser Ile Phe Phe Ala Asn Asn Arg Ser Tyr Thr Arg  
 85 90 95  
 Asp Ser Tyr Lys Met Ala Gly Met Ala Asp Asn Ile Glu Asp Leu Leu  
 100 105 110  
 His Phe Cys Arg Gln Met Phe Ser Met Lys Val Asp Asn Val Glu Tyr  
 115 120 125  
 Ala Leu Leu Thr Ala Ile Val Ile Phe Ser Asp Arg Pro Gly Leu Glu  
 130 135 140



## RH0020.ST25

Lys Ala Gln Leu Val Glu Ala Ile Gln Ser Tyr Tyr Ile Asp Thr Leu  
 145 150 155 160  
 Arg Ile Tyr Ile Leu Asn Arg His Cys Gly Asp Ser Met Ser Leu Val  
 165 170 175  
 Phe Tyr Ala Lys Leu Leu Ser Ile Leu Thr Glu Leu Arg Thr Leu Gly  
 180 185 190  
 Asn Gln Asn Ala Glu Met Cys Phe Ser Leu Lys Leu Lys Asn Arg Lys  
 195 200 205  
 Leu Pro Lys Phe Leu Glu Glu Ile Trp Asp Val His Ala Ile Pro Pro  
 210 215 220  
 Ser Val Gln Ser His Leu Gln Ile Thr Gln Glu Glu Asn Glu Arg Leu  
 225 230 235 240  
 Glu Arg Ala Glu Arg Met Arg Ala Ser Val Gly Gly Ala Ile Thr Ala  
 245 250 255  
 Gly Ile Asp Cys Asp Ser Ala Ser Thr Ser Ala Ala Ala Ala Ala Ala  
 260 265 270  
 Gln His Gln Pro Gln Pro Gln Pro Gln Pro Gln Pro Ser Ser Leu Thr  
 275 280 285  
 Gln Asn Asp Ser Gln His Gln Thr Gln Pro Gln Leu Gln Pro Gln Leu  
 290 295 300  
 Pro Pro Gln Leu Gln Gly Gln Leu Gln Pro Gln Leu Gln Pro Gln Leu  
 305 310 315 320  
 Gln Thr Gln Leu Gln Pro Gln Ile Gln Pro Gln Pro Gln Leu Leu Pro  
 325 330 335  
 Val Ser Ala Pro Val Pro Ala Ser Val Thr Ala Pro Gly Ser Leu Ser  
 340 345 350  
 Ala Val Ser Thr Ser Ser Glu Tyr Met Gly Gly Ser Ala Ala Ile Gly  
 355 360 365  
 Pro Ile Thr Pro Ala Thr Thr Ser Ser Ile Thr Ala Ala Val Thr Ala  
 370 375 380  
 Ser Ser Thr Thr Ser Ala Val Pro Met Gly Asn Gly Val Gly Val Gly  
 385 390 395 400  
 Val Gly Val Gly Gly Asn Val Ser Met Tyr Ala Asn Ala Gln Thr Ala  
 405 410 415  
 Met Ala Leu Met Gly Val Ala Leu His Ser His Gln Glu Gln Leu Ile  
 420 425 430  
 Gly Gly Val Ala Val Lys Ser Glu His Ser Thr Thr Ala  
 435 440 445

<210> 20  
 <211> 323  
 <212> PRT  
 <213> Artificial Sequence

<400> 20

Arg Pro Glu Cys Val Val Pro Glu Asn Gln Cys Ala Met Lys Arg Arg

[illegible]

23

RH0020.ST25

&lt;213&gt; Artificial Sequence

&lt;400&gt; 21

tggtgctatct	gtggggaccg	ctcctcaggc	aaacactatg	gggtatacag	ttgtgagggc	60
tgcaagggct	tcttcaagag	gacagtacgc	aaagacctga	cctacacctg	ccgagacaac	120
aaggactgcc	tgatcgacaa	gagacagcgg	aaccggtgtc	agtactgccg	ctaccagaag	180
tgcttgccca	tgggcatgaa	gcgggaagct	gtgcaggagg	agcggcagcg	gggcaaggac	240
cggaatgaga	acgaggtgga	gtccaccagc	agtgccaacg	aggacatgcc	tgtagagaag	300
attctggaag	ccgagcttgc	tgtcgagccc	aagactgaga	catacgtgga	ggcaaacatg	360
gggctgaacc	ccagctcacc	aaatgaccct	gttaccaaca	tctgtcaagc	agcagacaag	420
cagctcttca	ctcttgtgga	gtgggccaag	aggatccac	acttttctga	gctgccccta	480
gacgaccagg	tcattctgct	acgggcaggc	tggaacgagc	tgctgatcgc	ctccttctcc	540
caccgctcca	tagctgtgaa	agatgggatt	ctcctggcca	ccggcctgca	cgtacaccgg	600
aacagcgctc	acagtgtctg	ggtgggcgcc	atctttgaca	gggtgctaac	agagctgggtg	660
tctaagatgc	gtgacatgca	gatggacaag	acggagctgg	gctgcctgcg	agccattgtc	720
ctgttcaacc	ctgactctaa	ggggctctca	aaccctgctg	aggtggaggc	gttgagggag	780
aaggtgtatg	cgtcactaga	agcgtactgc	aaacacaagt	accctgagca	gccgggcagg	840
tttgccaagc	tgtgtctccg	cctgcctgca	ctgcgttcca	tggggctcaa	gtgcctggag	900
cacctgttct	tcttcaagct	catcggggac	acgcccacgc	acaccttcct	catggagatg	960
ctggaggcac	cacatcaagc	cacctag				987

&lt;210&gt; 22

&lt;211&gt; 789

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 22

aagcgggaag	ctgtgcagga	ggagcggcag	cggggcaagg	accggaatga	gaacgaggtg	60
gagtccacca	gcagtgccaa	cgaggacatg	cctgtagaga	agattctgga	agccgagctt	120
gctgtcgagc	ccaagactga	gacatacgtg	gaggcaaaca	tggggctgaa	ccccagctca	180
ccaaatgacc	ctgttaccaa	catctgtcaa	gcagcagaca	agcagctctt	cactcttgtg	240
gagtgggcca	agaggatccc	acacttttct	gagctgcccc	tagacgacca	ggteatcctg	300
ctacgggcag	gctggaacga	gctgtgatc	gcctccttct	cccaccgctc	catagctgtg	360
aaagatggga	ttctcctggc	caccggcctg	cacgtacacc	ggaacagcgc	tcacagtgtc	420
gggggtggcg	ccatctttga	cagggtgcta	acagagctgg	tgtctaagat	gcgtgacatg	480
cagatggaca	agacggagct	gggctgcctg	cgagccattg	tcctgttcaa	ccctgactct	540
aaggggctct	caaaccctgc	tgaggtggag	gcgttgaggg	agaaggtgta	tgcgctacta	600
gaagcgtact	gcaaacacaa	gtaccctgag	cagccgggca	ggtttgccaa	gctgctgctc	660

RH0020.ST25

cgctgcctg cactgcgttc catcgggctc aagtgcctgg agcacctgtt cttcttcaag	720
ctcatcgggg acacgcccac cgacaccttc ctcatggaga tgctggaggc accacatcaa	780
gccacctag	789

<210> 23  
 <211> 714  
 <212> DNA  
 <213> Artificial Sequence

<400> 23	
gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag	60
actgagacat acgtggaggc aaacatgggg ctgaaccca gctcaccaa tgacctgtt	120
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg	180
atcccacact tttctgagct gcccctagac gaccaggtca tcctgctacg ggcaggctgg	240
aacgagctgc tgatgcctc cttctccac cgctccatag ctgtgaaaga tgggattctc	300
ctggccaccg gcctgcacgt acaccggaac agcgctcaca gtgctggggg gggcgccatc	360
tttgacaggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg	420
gagctgggct gcctgcgagc cattgtcctg ttcaacctg actctaagg gctctcaaac	480
cctgctgagg tggaggcggt gagggagaag gtgtatgct cactagaagc gtactgcaaa	540
cacaagtacc ctgagcagcc gggcaggttt gccaaagctgc tgctccgct gcctgcactg	600
cgttccatcg ggctcaagt cctggagcac ctgttcttct tcaagctcat cggggacacg	660
cccatcgaca ccttctcat ggagatgctg gaggcaccac atcaagccac ctag	714

<210> 24  
 <211> 536  
 <212> DNA  
 <213> Artificial Sequence

<400> 24	
ggatcccaca cttttctgag ctgcccctag acgaccaggt catcctgcta cgggcaggct	60
ggaacgagct gctgatcgcc tccttctccc accgctccat agctgtgaaa gatgggattc	120
tcctggccac cggcctgcac gtacaccgga acagcgctca cagtgcctgg gtggcgcca	180
tccttgacag ggtgctaaca gagctggtgt ctaagatgcg tgacatgcag atggacaaga	240
cggagctggg ctgcctgcga gccattgtcc tgttcaacct tgactctaag gggctctcaa	300
accctgctga ggtggaggcg ttgagggaga aggtgtatgc gtcactagaa gcgtactgca	360
aacacaagta ccctgagcag ccgggcagg ttgccaagct gctgctccgc ctgcctgcac	420
tgcgttccat cgggctcaag tgctggagc acctgttctt cttcaagctc atcggggaca	480
cggccatcga caccttctc atggagatgc tggaggcacc acatcaagcc acctag	536

<210> 25

RH0020.ST25

&lt;211&gt; 672

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 25

```

gccaacgagg acatgcctgt agagaagatt ctggaagccg agcttgctgt cgagcccaag      60
actgagacat acgtggaggc aaacatgggg ctgaaccca gctcaccaaa tgaccctgtt      120
accaacatct gtcaagcagc agacaagcag ctcttcactc ttgtggagtg ggccaagagg      180
atcccacact tttctgagct gcccctagac gaccaggtca tcctgctacg ggcaggctgg      240
aacgagctgc tgatgcctc cttctccac cgctccatag ctgtgaaaga tgggattctc      300
ctggccaccg gcctgcacgt acaccgaac agcgctcaca gtgctggggt gggcgccatc      360
tttgacaggg tgctaacaga gctggtgtct aagatgcgtg acatgcagat ggacaagacg      420
gagctgggct gcctgcgagc cattgtcctg ttcaaccctg actctaagg gctctcaaac      480
cctgctgagg tggaggcgtt gagggagaag gtgtatgcgt cactagaagc gtactgcaaa      540
cacaagtacc ctgagcagcc gggcaggttt gccaagctgc tgctccgcct gcctgcactg      600
cgttccatcg ggctcaagtg cctggagcac ctgttcttct tcaagctcat cggggacacg      660
cccatcgaca cc                                     672

```

&lt;210&gt; 26

&lt;211&gt; 1123

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;400&gt; 26

```

tgcgccatct gcggggaccg ctctcaggc aagcactatg gagtgtacag ctgcgagggg      60
tgcaagggct tcttcaagcg gacggtgcgc aaggacctga cctacacctg ccgcgacaac      120
aaggactgcc tgattgacaa gcggcagcgg aaccggtgcc agtactgccg ctaccagaag      180
tgcttgccca tgggcatgaa gcgggaagcc gtgcaggagg agcggcagcg tggcaaggac      240
cggaacgaga atgaggtgga gtcgaccagc agcgccaacg aggacatgcc ggtggagagg      300
atcctggagg ctgagctggc cgtggagccc aagaccgaga cctacgtgga ggcaaacatg      360
gggctgaacc ccagctcgcc gaacgacctt gtcaccaaca ttgccaagc agccgacaaa      420
cagcttttca ccctggtgga gtgggccaag cggatccac acttctcaga gctgcccctg      480
gacgaccagg tcactctgct gcgggcaggc tggaatgagc tgctcatcgc ctcttctcc      540
caccgctcca tcgccgtgaa ggacgggatc ctctggcca ccgggctgca cgtccaccgg      600
aacagcgccc acagcgcagg ggtgggcgcc atctttgaca ggggtgctgac ggagcttgtg      660
tccaagatgc gggacatgca gatggacaag acggagctgg gctgcctgcg cgccatcgtc      720
ctctttaacc ctgactccaa ggggctctcg aaccggccg aggtggaggc gctgaggagg      780
aaggctctat cgctccttga ggcctactgc aagcacaagt acccagagca gccgggaagg      840
ttcgctaagc tcttgctccg cctgccggct ctgcgctcca tcgggctcaa atgcctggaa      900

```

RH0020.ST25

```

catctcttct ttttcaagct catcggggac acacccattg acaccttcct tatggagatg   960
ctggaggcgc cgcaccaa at gacttaggcc tgcgggcca tcttttgtgc ccaccgttc   1020
tggccaccct gcttgagcgc cagctgttct tctcagcctg agccctgtcc ctgccttct   1080
ctgcctggcc tgtttggact ttggggcaca gcctgtcact gct                       1123

```

<210> 27  
 <211> 925  
 <212> DNA  
 <213> Artificial Sequence

```

<400> 27
aagcgggaag ccgtgcagga ggagcggcag cgtggcaagg accggaacga gaatgaggtg   60
gagtcgacca gcagcgccaa cgaggacatg ccggtggaga ggatcctgga ggctgagctg   120
gccgtggagc ccaagaccga gacctacgtg gaggcaaaca tggggctgaa cccagctcg   180
ccgaacgacc ctgtcaccaa cat ttgcca gacagccaca aacagctttt caccctgggtg   240
gagtgggcca agcggatccc acacttctca gagctgcccc tggacgacca ggtcatcctg   300
ctgcgggcag gctggaatga gctgtcctc gcctccttct cccaccgtc catcgccgtg   360
aaggacggga tcctcctggc caccgggctg cagctccacc ggaacagcgc ccacagcgca   420
gggggtggcg ccatctttga caggtgtctg acggagcttg tgtccaagat gcgggacatg   480
cagatggaca agacggagct gggctgcctg cgcgccatcg tcctctttaa ccctgactcc   540
aaggggctct cgaaccggc cgaggtggag gcgctgaggg agaaggtcta tgcgtccttg   600
gaggcctact gcaagcaca gtacccagag cagccgggaa ggttcgctaa gctcttgctc   660
cgcctgccg ctctgcgctc catcgggctc aaatgcctgg aacatctctt cttcttcaag   720
ctcatcgggg acacacccat tgacaccttc cttatggaga tgctggaggc gccgcaccaa   780
atgacttagg cctgcggggc catcctttgt gccaccctg tctggccacc ctgcctggac   840
gccagctgtt cttctcagcc tgagccctgt ccctgccctt ctctgcctgg cctgtttgga   900
ctttggggca cagcctgtca ctgct                                     925

```

<210> 28  
 <211> 850  
 <212> DNA  
 <213> Artificial Sequence

```

<400> 28
gccaacgagg acatgccggt ggagaggatc ctggaggctg agctggccgt ggagcccaag   60
accgagacct acgtggaggc aaacatgggg ctgaaccca gctcgccgaa cgaccctgtc   120
accaacattt gccaaagcgc cgacaaacag cttttcacc tgggtggagt ggccaagcgg   180
atccacact tctcagagct gcccctggac gaccaggtca tcctgctgcg ggcaggctgg   240
aatgagctgc tcatgcctc cttctccac cgctccatcg ccgtgaagga cgggatcctc   300

```

RH0020.ST25

```

ctggccaccg ggctgcacgt ccaccggaac agcgcccaca ggcgaggggt gggcgccatc 360
tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 420
gagctgggct gcctgcgcgc catcgtcctc tttaaccctg actccaaggg gctctcgaac 480
ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgaggc ctactgcaag 540
cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggctctg 600
cgctccatcg ggctcaaagt cctggaacat ctcttcttct tcaagctcat cggggacaca 660
cccattgaca ccttccttat ggagatgctg gaggcgccgc accaaatgac ttaggcctgc 720
gggcccattc tttgtgcca ccggttctg ccaccctgcc tggacgccag ctgttcttct 780
cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 840
tgctactgct 850

```

<210> 29  
<211> 670  
<212> DNA  
<213> Artificial Sequence

```

<400> 29
atcccacact tctcagagct gcccctggac gaccaggcca tctgctgctg ggcaggctgg 60
aatgagctgc tcatgcctc cttctccac cgctccatcg ccgtgaagga cgggatcctc 120
ctggccaccg ggctgcacgt ccaccggaac agcgcccaca ggcgaggggt gggcgccatc 180
tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 240
gagctgggct gcctgcgcgc catcgtcctc tttaaccctg actccaaggg gctctcgaac 300
ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttgaggc ctactgcaag 360
cacaagtacc cagagcagcc gggaagggtc gctaagctct tgctccgcct gccggctctg 420
cgctccatcg ggctcaaagt cctggaacat ctcttcttct tcaagctcat cggggacaca 480
cccattgaca ccttccttat ggagatgctg gaggcgccgc accaaatgac ttaggcctgc 540
gggcccattc tttgtgcca ccggttctg ccaccctgcc tggacgccag ctgttcttct 600
cagcctgagc cctgtccctg cccttctctg cctggcctgt ttggactttg gggcacagcc 660
tgctactgct 670

```

<210> 30  
<211> 672  
<212> DNA  
<213> Artificial Sequence

```

<400> 30
gccaacgagg acatgccggt ggagaggatc ctggaggctg agctggccgt ggagcccaag 60
accgagacct acgtggaggc aaacatgggg ctgaaccca gctcgccgaa cgaccctgtc 120
accaacattt gccaagcagc cgacaacag cttttcacc ttggtggagt ggccaagcgg 180
atcccacact tctcagagct gcccctggac gaccaggcca tctgctgctg ggcaggctgg 240

```

RH0020.ST25

```

aatgagctgc tcatcgctc cttctccac cgctccatcg ccgtgaagga cgggacctc 300
ctggccaccg ggctgcacgt ccaccggaac agcgcccaca gcgcaggggt gggcgccatc 360
tttgacaggg tgctgacgga gcttgtgtcc aagatgcggg acatgcagat ggacaagacg 420
gagctgggct gcctgcgcgc catcgtcctc ttttaaccctg actccaagg gctctcgaac 480
ccggccgagg tggaggcgct gagggagaag gtctatgcgt ccttggaggc ctactgcaag 540
cacaagtacc cagagcagcc gggaaggttc gctaagctct tgctccgcct gccggctctg 600
cgctccatcg ggctcaaagt cctggaacat ctcttcttct tcaagctcat cggggacaca 660
cccattgaca cc 672

```

```

<210> 31
<211> 328
<212> PRT
<213> Artificial Sequence

```

```

<400> 31

```

```

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr
1      5      10
Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp
20     25     30
Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg
35     40     45
Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met
50     55     60
Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp
65     70     75     80
Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met
85     90     95
Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr
100    105    110
Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn
115    120    125
Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr
130    135    140
Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu
145    150    155    160
Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile
165    170    175
Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu
180    185    190
Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val
195    200    205
Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg

```



210				215				RH0020.ST25 220							
Asp 225	Met	Gln	Met	Asp	Lys 230	Thr	Glu	Leu	Gly	Cys 235	Leu	Arg	Ala	Ile	Val 240
Leu	Phe	Asn	Pro	Asp 245	Ser	Lys	Gly	Leu	Ser 250	Asn	Pro	Ala	Glu	Val 255	Glu
Ala	Leu	Arg	Glu 260	Lys	Val	Tyr	Ala	Ser 265	Leu	Glu	Ala	Tyr	Cys 270	Lys	His
Lys	Tyr	Pro 275	Glu	Gln	Pro	Gly	Arg 280	Phe	Ala	Lys	Leu	Leu 285	Leu	Arg	Leu
Pro	Ala 290	Leu	Arg	Ser	Ile	Gly 295	Leu	Lys	Cys	Leu	Glu 300	His	Leu	Phe	Phe
Phe 305	Lys	Leu	Ile	Gly	Asp 310	Thr	Pro	Ile	Asp	Thr 315	Phe	Leu	Met	Glu	Met 320
Leu	Glu	Ala	Pro	His 325	Gln	Ala	Thr								
<210> 32															
<211> 262															
<212> PRT															
<213> Artificial Sequence															
<400> 32															
Lys 1	Arg	Glu	Ala	Val 5	Gln	Glu	Glu	Arg	Gln 10	Arg	Gly	Lys	Asp	Arg 15	Asn
Glu	Asn	Glu	Val 20	Glu	Ser	Thr	Ser	Ser 25	Ala	Asn	Glu	Asp	Met 30	Pro	Val
Glu	Lys 35	Ile	Leu	Glu	Ala	Glu	Leu 40	Ala	Val	Glu	Pro	Lys 45	Thr	Glu	Thr
Tyr	Val 50	Glu	Ala	Asn	Met	Gly 55	Leu	Asn	Pro	Ser	Ser 60	Pro	Asn	Asp	Pro
Val 65	Thr	Asn	Ile	Cys	Gln 70	Ala	Ala	Asp	Lys	Gln 75	Leu	Phe	Thr	Leu	Val 80
Glu	Trp	Ala	Lys 85	Arg	Ile	Pro	His	Phe	Ser 90	Glu	Leu	Pro	Leu	Asp 95	Asp
Gln	Val	Ile 100	Leu	Leu	Arg	Ala	Gly	Trp 105	Asn	Glu	Leu	Leu	Ile 110	Ala	Ser
Phe	Ser 115	His	Arg	Ser	Ile	Ala 120	Val	Lys	Asp	Gly	Ile	Leu 125	Leu	Ala	Thr
Gly 130	Leu	His	Val	His	Arg	Asn 135	Ser	Ala	His	Ser	Ala 140	Gly	Val	Gly	Ala
Ile 145	Phe	Asp	Arg	Val	Leu 150	Thr	Glu	Leu	Val	Ser 155	Lys	Met	Arg	Asp	Met 160
Gln	Met	Asp	Lys 165	Thr	Glu	Leu	Gly	Cys	Leu 170	Arg	Ala	Ile	Val	Leu 175	Phe
Asn	Pro	Asp 180	Ser	Lys	Gly	Leu	Ser	Asn 185	Pro	Ala	Glu	Val	Glu 190	Ala	Leu

RH0020.ST25

Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr  
 195 200 205  
 Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Arg Leu Pro Ala  
 210 215 220  
 Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys  
 225 230 235 240  
 Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu  
 245 250 255  
 Ala Pro His Gln Ala Thr  
 260

<210> 33  
 <211> 237  
 <212> PRT  
 <213> Artificial Sequence

<400> 33

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala  
 1 5 10 15  
 Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
 20 25 30  
 Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
 35 40 45  
 Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
 50 55 60  
 Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
 65 70 75 80  
 Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
 85 90 95  
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
 100 105 110  
 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
 115 120 125  
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
 130 135 140  
 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
 145 150 155 160  
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
 165 170 175  
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
 180 185 190  
 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
 195 200 205  
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
 210 215 220

RH0020.ST25

Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala Thr  
 225 230 235

<210> 34  
 <211> 177  
 <212> PRT  
 <213> Artificial Sequence

<400> 34

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu  
 1 5 10 15

Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser  
 20 25 30

Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His  
 35 40 45

Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val  
 50 55 60

Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr  
 65 70 75 80

Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys  
 85 90 95

Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr  
 100 105 110

Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly  
 115 120 125

Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly  
 130 135 140

Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr  
 145 150 155 160

Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Ala  
 165 170 175

Thr

<210> 35  
 <211> 224  
 <212> PRT  
 <213> Artificial Sequence

<400> 35

Ala Asn Glu Asp Met Pro Val Glu Lys Ile Leu Glu Ala Glu Leu Ala  
 1 5 10 15

Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
 20 25 30

Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
 35 40 45

Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
 50 55 60

## RH0020.ST25

Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
 65 70 75 80  
 Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
 85 90 95  
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
 100 105 110  
 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
 115 120 125  
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
 130 135 140  
 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
 145 150 155 160  
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
 165 170 175  
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
 180 185 190  
 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
 195 200 205  
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
 210 215 220  
 <210> 36  
 <211> 328  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 36

Cys Ala Ile Cys Gly Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr  
 1 5 10 15  
 Ser Cys Glu Gly Cys Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp  
 20 25 30  
 Leu Thr Tyr Thr Cys Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg  
 35 40 45  
 Gln Arg Asn Arg Cys Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met  
 50 55 60  
 Gly Met Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp  
 65 70 75 80  
 Arg Asn Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met  
 85 90 95  
 Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr  
 100 105 110  
 Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn  
 115 120 125

RH0020.ST25

Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr  
 130 135 140  
 Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu  
 145 150 155 160  
 Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile  
 165 170 175  
 Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu  
 180 185 190  
 Ala Thr Gly Leu His Val His Arg Asn Ser Ala His Ser Ala Gly Val  
 195 200 205  
 Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg  
 210 215 220  
 Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val  
 225 230 235 240  
 Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu  
 245 250 255  
 Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His  
 260 265 270  
 Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu  
 275 280 285  
 Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe  
 290 295 300  
 Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met  
 305 310 315 320  
 Leu Glu Ala Pro His Gln Met Thr  
 325

<210> 37  
 <211> 262  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 37

Lys Arg Glu Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn  
 1 5 10 15  
 Glu Asn Glu Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val  
 20 25 30  
 Glu Arg Ile Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr  
 35 40 45  
 Tyr Val Glu Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro  
 50 55 60  
 Val Thr Asn Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val

RH0020.ST25

65		70		75		80									
Glu	Trp	Ala	Lys	Arg	Ile	Pro	His	Phe	Ser	Glu	Leu	Pro	Leu	Asp	Asp
			85						90					95	
Gln	Val	Ile	Leu	Leu	Arg	Ala	Gly	Trp	Asn	Glu	Leu	Leu	Ile	Ala	Ser
			100					105					110		
Phe	Ser	His	Arg	Ser	Ile	Ala	Val	Lys	Asp	Gly	Ile	Leu	Leu	Ala	Thr
		115					120					125			
Gly	Leu	His	Val	His	Arg	Asn	Ser	Ala	His	Ser	Ala	Gly	Val	Gly	Ala
		130				135					140				
Ile	Phe	Asp	Arg	Val	Leu	Thr	Glu	Leu	Val	Ser	Lys	Met	Arg	Asp	Met
145					150					155					160
Gln	Met	Asp	Lys	Thr	Glu	Leu	Gly	Cys	Leu	Arg	Ala	Ile	Val	Leu	Phe
			165						170					175	
Asn	Pro	Asp	Ser	Lys	Gly	Leu	Ser	Asn	Pro	Ala	Glu	Val	Glu	Ala	Leu
			180					185					190		
Arg	Glu	Lys	Val	Tyr	Ala	Ser	Leu	Glu	Ala	Tyr	Cys	Lys	His	Lys	Tyr
		195					200					205			
Pro	Glu	Gln	Pro	Gly	Arg	Phe	Ala	Lys	Leu	Leu	Leu	Arg	Leu	Pro	Ala
		210				215					220				
Leu	Arg	Ser	Ile	Gly	Leu	Lys	Cys	Leu	Glu	His	Leu	Phe	Phe	Phe	Lys
225					230					235					240
Leu	Ile	Gly	Asp	Thr	Pro	Ile	Asp	Thr	Phe	Leu	Met	Glu	Met	Leu	Glu
			245						250					255	
Ala	Pro	His	Gln	Met	Thr										
			260												

<210> 38  
 <211> 237  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 38

Ala	Asn	Glu	Asp	Met	Pro	Val	Glu	Arg	Ile	Leu	Glu	Ala	Glu	Leu	Ala
1				5					10					15	
Val	Glu	Pro	Lys	Thr	Glu	Thr	Tyr	Val	Glu	Ala	Asn	Met	Gly	Leu	Asn
			20					25					30		
Pro	Ser	Ser	Pro	Asn	Asp	Pro	Val	Thr	Asn	Ile	Cys	Gln	Ala	Ala	Asp
		35					40					45			
Lys	Gln	Leu	Phe	Thr	Leu	Val	Glu	Trp	Ala	Lys	Arg	Ile	Pro	His	Phe
	50				55				60						
Ser	Glu	Leu	Pro	Leu	Asp	Asp	Gln	Val	Ile	Leu	Leu	Arg	Ala	Gly	Trp
65					70					75					80

## RH0020.ST25

Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
 85 90 95  
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
 100 105 110  
 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
 115 120 125  
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
 130 135 140  
 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
 145 150 155 160  
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
 165 170 175  
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
 180 185 190  
 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
 195 200 205  
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr  
 210 215 220  
 Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met Thr  
 225 230 235

<210> 39  
 <211> 177  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 39

Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu  
 1 5 10 15  
 Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser  
 20 25 30  
 Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His  
 35 40 45  
 Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val  
 50 55 60  
 Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr  
 65 70 75 80  
 Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys  
 85 90 95  
 Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr  
 100 105 110  
 Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly  
 115 120 125

RH0020.ST25

Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly  
 130 135 140  
 Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr  
 145 150 155 160  
 Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His Gln Met  
 165 170 175

Thr

<210> 40  
 <211> 224  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

&lt;400&gt; 40

Ala Asn Glu Asp Met Pro Val Glu Arg Ile Leu Glu Ala Glu Leu Ala  
 1 5 10 15  
 Val Glu Pro Lys Thr Glu Thr Tyr Val Glu Ala Asn Met Gly Leu Asn  
 20 25 30  
 Pro Ser Ser Pro Asn Asp Pro Val Thr Asn Ile Cys Gln Ala Ala Asp  
 35 40 45  
 Lys Gln Leu Phe Thr Leu Val Glu Trp Ala Lys Arg Ile Pro His Phe  
 50 55 60  
 Ser Glu Leu Pro Leu Asp Asp Gln Val Ile Leu Leu Arg Ala Gly Trp  
 65 70 75 80  
 Asn Glu Leu Leu Ile Ala Ser Phe Ser His Arg Ser Ile Ala Val Lys  
 85 90 95  
 Asp Gly Ile Leu Leu Ala Thr Gly Leu His Val His Arg Asn Ser Ala  
 100 105 110  
 His Ser Ala Gly Val Gly Ala Ile Phe Asp Arg Val Leu Thr Glu Leu  
 115 120 125  
 Val Ser Lys Met Arg Asp Met Gln Met Asp Lys Thr Glu Leu Gly Cys  
 130 135 140  
 Leu Arg Ala Ile Val Leu Phe Asn Pro Asp Ser Lys Gly Leu Ser Asn  
 145 150 155 160  
 Pro Ala Glu Val Glu Ala Leu Arg Glu Lys Val Tyr Ala Ser Leu Glu  
 165 170 175  
 Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln Pro Gly Arg Phe Ala Lys  
 180 185 190  
 Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile Gly Leu Lys Cys Leu  
 195 200 205  
 Glu His Leu Phe Phe Phe Lys Leu Ile Gly Asp Thr Pro Ile Asp Thr



RH0020.ST25

210

215

220

<210> 41  
 <211> 441  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 41  
 atgaagctac tgtcttctat cgaacaagca tgcgatattt gccgacttaa aaagctcaag 60  
 tgctccaaag aaaaaccgaa gtgcgccaag tgtctgaaga acaactggga gtgtcgctac 120  
 tctcccaaaa caaaagggtc tccgctgact agggcacatc tgacagaagt ggaatcaagg 180  
 ctagaaagac tggaacagct atttctactg atttttcctc gagaagacct tgacatgatt 240  
 ttgaaaatgg attctttaca ggatataaaa gcattgttaa caggattatt tgtacaagat 300  
 aatgtgaata aagatgccgt cacagataga ttggcttcag tggagactga tatgcctcta 360  
 acattgagac agcatagaat aagtgcgaca tcatcatcgg aagagagtag taacaaagg 420  
 caagacagt tgactgtatc g 441

<210> 42  
 <211> 147  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 42

Met	Lys	Leu	Leu	Ser	Ser	Ile	Glu	Gln	Ala	Cys	Asp	Ile	Cys	Arg	Leu
1				5					10					15	
Lys	Lys	Leu	Lys	Cys	Ser	Lys	Glu	Lys	Pro	Lys	Cys	Ala	Lys	Cys	Leu
			20					25					30		
Lys	Asn	Asn	Trp	Glu	Cys	Arg	Tyr	Ser	Pro	Lys	Thr	Lys	Arg	Ser	Pro
		35					40					45			
Leu	Thr	Arg	Ala	His	Leu	Thr	Glu	Val	Glu	Ser	Arg	Leu	Glu	Arg	Leu
	50					55					60				
Glu	Gln	Leu	Phe	Leu	Leu	Ile	Phe	Pro	Arg	Glu	Asp	Leu	Asp	Met	Ile
65					70					75				80	
Leu	Lys	Met	Asp	Ser	Leu	Gln	Asp	Ile	Lys	Ala	Leu	Leu	Thr	Gly	Leu
				85					90					95	
Phe	Val	Gln	Asp	Asn	Val	Asn	Lys	Asp	Ala	Val	Thr	Asp	Arg	Leu	Ala
			100					105					110		
Ser	Val	Glu	Thr	Asp	Met	Pro	Leu	Thr	Leu	Arg	Gln	His	Arg	Ile	Ser
		115					120					125			

RH0020.ST25

Ala Thr Ser Ser Ser Glu Glu Ser Ser Asn Lys Gly Gln Arg Gln Leu  
 130 135 140

Thr Val Ser  
 145

<210> 43  
 <211> 606  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 43  
 atgaaagcgt taacggccag gcaacaagag gtgtttgatc tcatccgtga tcacatcagc 60  
 cagacaggtg tgccgccgac gcgtgcggaa atcgcgccagc gtttgggggtt ccgttcccca 120  
 aacgcggctg aagaacatct gaaggcgtg gcacgcaaag gcgttattga aattgtttcc 180  
 ggcgcacac gcgggattcg tctgttcag gaagaggaag aagggttgcc gctggtaggt 240  
 cgtgtggctg ccggtgaacc acttctggcg caacagcata ttgaaggtca ttatcaggtc 300  
 gatccttcct tattcaagcc gaatgctgat ttctgtctgc gcgtcagcgg gatgtcgatg 360  
 aaagatatcg gcattatgga tggtagcttg ctggcagtcg ataaaactca ggatgtacgt 420  
 aacggtcagg tcgttgtcgc acgtattgat gacgaagtta ccgttaagcg cctgaaaaaa 480  
 cagggcaata aagtcgaact gttgccagaa aatagcgagt ttaaaccaat tgtcgtagat 540  
 cttcgtcagc agagcttcac cattgaaggg ctggcggttg gggttattcg caacggcgac 600  
 tggctg 606

<210> 44  
 <211> 202  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 44

Met Lys Ala Leu Thr Ala Arg Gln Gln Glu Val Phe Asp Leu Ile Arg  
 1 5 10 15  
 Asp His Ile Ser Gln Thr Gly Met Pro Pro Thr Arg Ala Glu Ile Ala  
 20 25 30  
 Gln Arg Leu Gly Phe Arg Ser Pro Asn Ala Ala Glu Glu His Leu Lys  
 35 40 45  
 Ala Leu Ala Arg Lys Gly Val Ile Glu Ile Val Ser Gly Ala Ser Arg  
 50 55 60

RH0020.ST25

Gly Ile Arg Leu Leu Gln Glu Glu Glu Glu Gly Leu Pro Leu Val Gly  
 65 70 75 80  
 Arg Val Ala Ala Gly Glu Pro Leu Leu Ala Gln Gln His Ile Glu Gly  
 85 90 95  
 His Tyr Gln Val Asp Pro Ser Leu Phe Lys Pro Asn Ala Asp Phe Leu  
 100 105 110  
 Leu Arg Val Ser Gly Met Ser Met Lys Asp Ile Gly Ile Met Asp Gly  
 115 120 125  
 Asp Leu Leu Ala Val His Lys Thr Gln Asp Val Arg Asn Gly Gln Val  
 130 135 140  
 Val Val Ala Arg Ile Asp Asp Glu Val Thr Val Lys Arg Leu Lys Lys  
 145 150 155 160  
 Gln Gly Asn Lys Val Glu Leu Leu Pro Glu Asn Ser Glu Phe Lys Pro  
 165 170 175  
 Ile Val Val Asp Leu Arg Gln Gln Ser Phe Thr Ile Glu Gly Leu Ala  
 180 185 190  
 Val Gly Val Ile Arg Asn Gly Asp Trp Leu  
 195 200

<210> 45  
 <211> 271  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 45  
 atgggcccta aaaagaagcg taaagtcgcc cccccgaccg atgtcagcct gggggacgag 60  
 ctccacttag acggcgagga cgtggcgatg ggcgatgccg acgcgctaga cgatttcgat 120  
 ctggacatgt tgggggacgg ggattccccg gggccgggat ttacccccca cgactccgcc 180  
 ccctacggcg ctctggatat ggccgacttc gagtttgagc agatgtttac cgatgccctt 240  
 ggaattgacg agtacggtgg ggaattccccg g 271

<210> 46  
 <211> 90  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 46

Met Gly Pro Lys Lys Lys Arg Lys Val Ala Pro Pro Thr Asp Val Ser  
 1 5 10 15  
 Leu Gly Asp Glu Leu His Leu Asp Gly Glu Asp Val Ala Met Ala His

	20	25	RH0020.ST25	30
Ala Asp Ala Leu Asp Asp Phe Asp Leu Asp Met Leu Gly Asp Gly Asp	35	40		45
Ser Pro Gly Pro Gly Phe Thr Pro His Asp Ser Ala Pro Tyr Gly Ala	50	55		60
Leu Asp Met Ala Asp Phe Glu Phe Glu Gln Met Phe Thr Asp Ala Leu	65	70	75	80
Gly Ile Asp Glu Tyr Gly Gly Glu Phe Pro	85	90		
<210>	47			
<211>	19			
<212>	DNA			
<213>	Artificial Sequence			
<220>				
<221>	misc_feature			
<223>	Novel Sequence			
<400>	47			
ggagtactgt cctccgagc				19
<210>	48			
<211>	666			
<212>	DNA			
<213>	Artificial Sequence			
<220>				
<221>	misc_feature			
<223>	Novel Sequence			
<400>	48			
ggatccccag cttggaattc gacaggttat cagcaacaac acagtcatat ccattctcaa				60
ttagctctac cacagtgtgt gaaccaatgt atccagcacc acctgtaacc aaaacaattt				120
tagaagtact ttcactttgt aactgagctg tcattttatat tgaattttca aaaattctta				180
cttttttttt ggatggacgc aaagaagttt aataatcata ttacatggca ttaccaccat				240
atacatatcc atatacatat ccatatctaa tcttacctcg actgctgtat ataaaaccag				300
tggttatatg tacagtactg ctgtatataa aaccagtggt tatatgtaca gtacgtcgac				360
tgctgtatat aaaaccagtg gttatatgta cagtaactgct gtatataaaa ccagtggtta				420
tatgtacagt acgtcgaggg atgataatgc gattagtttt ttagccttat ttctggggta				480
attaatcagc gaagcgatga tttttgatct attaacagat atataaatgc aaaaactgca				540
taaccacttt aactaatact ttcaacattht tctggtttgta ttacttctta ttcaaatgta				600
ataaaaagtat caacaaaaaa ttgttaatat acctctatac tttaacgtca aggagaaaaa				660
actata				666
<210>	49			

RH0020.ST25

<211> 1542  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 49  
 ctggacctga aacacgaagt ggcttaccga ggggtgctcc caggccaggt gaaggccgaa 60  
 ccgggggtcc acaacggcca ggtcaacggc cacgtgaggg actggatggc aggcggcgct 120  
 ggtgccaat cgcggtctcc gggagcgggtg gctcaacccc agcctaacaa tgggtattcg 180  
 tcgccactct cctcgggaag ctacgggccc tacagtccaa atgggaaaat aggcggtgag 240  
 gaactgtcgc cagcttcaag tataaatggg tgcagtacag atggcgaggc acgacgtcag 300  
 aagaagggcc ctgcgccccg tcagcaagag gaactgtgtc tggatatcggt ggacagagcc 360  
 tccggatacc actacaatgc gctcacgtgt gaagggtgta aagggttctt cagacggagt 420  
 gttaccaaaa atgcggttta ttttgtaaa ttccggtcacg cttgcgaaat ggacatgtac 480  
 atgcgacgga aatgccagga gtgccgcctg aagaagtgtc tagctgtagg catgaggcct 540  
 gagtgcgtag tacccgagac tcagtgcgcc atgaagcgga aagagaagaa agcacagaag 600  
 gagaaggaca aactgcctgt cagcacgacg acggtggacg accacatgcc gccattatg 660  
 cagtgtgaac ctccacctcc tgaagcagca aggattcacg aagtgggtccc aaggtttctc 720  
 tccgacaagc tgttgagac aaaccggcag aaaaacatcc ccagttgac agccaaccag 780  
 cagttcctta tcgccaggct catctggtac caggacgggt acgagcagcc ttctgatgaa 840  
 gatttgaaga ggattacgca gacgtggcag caagcggacg atgaaaacga agagtctgac 900  
 actcccttcc gccagatcac agagatgact atcctcacgg tccaacttat cgtggagttc 960  
 gcgaagggat tgccagggtt cgccaagatc tcgcagcctg atcaaattac gctgcttaag 1020  
 gcttgctcaa gtgaggtaat gatgtccga gtcgcgcgac gatacgatgc ggctcagac 1080  
 agtgttctgt tcgcgaacaa ccaagcgtac actcgcgaca actaccgcaa ggctggcatg 1140  
 gcctacgtca tcgaggatct actgcacttc tgccggtgca tgtactctat ggcgttggac 1200  
 aacatccatt acgcgtgtct cacggtgtc gtcactcttt ctgaccggcc agggttggag 1260  
 cagccgcaac tggtggaaga aatccagcgg tactacctga atacgtccg catctatatc 1320  
 ctgaaccagc tgagcgggtc ggcggttcg tccgtcatat acggcaagat cctctcaatc 1380  
 ctctctgagc tacgcacgct cggcatgcaa aactccaaca tgtgcatctc cctcaagctc 1440  
 aagaacagaa agctgccgcc tttcctcgag gagatctggg atgtggcgga catgtcgcac 1500  
 acccaaccgc cgctatcct cgagtcccc acgaatctct ag 1542

<210> 50  
 <211> 513

RH0020.ST25

<212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 50

Leu	Asp	Leu	Lys	His	Glu	Val	Ala	Tyr	Arg	Gly	Val	Leu	Pro	Gly	Gln	1	5	10	15
Val	Lys	Ala	Glu	Pro	Gly	Val	His	Asn	Gly	Gln	Val	Asn	Gly	His	Val	20	25	30	
Arg	Asp	Trp	Met	Ala	Gly	Gly	Ala	Gly	Ala	Asn	Ser	Pro	Ser	Pro	Gly	35	40	45	
Ala	Val	Ala	Gln	Pro	Gln	Pro	Asn	Asn	Gly	Tyr	Ser	Ser	Pro	Leu	Ser	50	55	60	
Ser	Gly	Ser	Tyr	Gly	Pro	Tyr	Ser	Pro	Asn	Gly	Lys	Ile	Gly	Arg	Glu	65	70	75	80
Glu	Leu	Ser	Pro	Ala	Ser	Ser	Ile	Asn	Gly	Cys	Ser	Thr	Asp	Gly	Glu	85	90	95	
Ala	Arg	Arg	Gln	Lys	Lys	Gly	Pro	Ala	Pro	Arg	Gln	Gln	Glu	Glu	Leu	100	105	110	
Cys	Leu	Val	Cys	Gly	Asp	Arg	Ala	Ser	Gly	Tyr	His	Tyr	Asn	Ala	Leu	115	120	125	
Thr	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe	Arg	Arg	Ser	Val	Thr	Lys	Asn	130	135	140	
Ala	Val	Tyr	Ile	Cys	Lys	Phe	Gly	His	Ala	Cys	Glu	Met	Asp	Met	Tyr	145	150	155	160
Met	Arg	Arg	Lys	Cys	Gln	Glu	Cys	Arg	Leu	Lys	Lys	Cys	Leu	Ala	Val	165	170	175	
Gly	Met	Arg	Pro	Glu	Cys	Val	Val	Pro	Glu	Thr	Gln	Cys	Ala	Met	Lys	180	185	190	
Arg	Lys	Glu	Lys	Lys	Ala	Gln	Lys	Glu	Lys	Asp	Lys	Leu	Pro	Val	Ser	195	200	205	
Thr	Thr	Thr	Val	Asp	Asp	His	Met	Pro	Pro	Ile	Met	Gln	Cys	Glu	Pro	210	215	220	
Pro	Pro	Pro	Glu	Ala	Ala	Arg	Ile	His	Glu	Val	Val	Pro	Arg	Phe	Leu	225	230	235	240
Ser	Asp	Lys	Leu	Leu	Glu	Thr	Asn	Arg	Gln	Lys	Asn	Ile	Pro	Gln	Leu	245	250	255	
Thr	Ala	Asn	Gln	Gln	Phe	Leu	Ile	Ala	Arg	Leu	Ile	Trp	Tyr	Gln	Asp	260	265	270	
Gly	Tyr	Glu	Gln	Pro	Ser	Asp	Glu	Asp	Leu	Lys	Arg	Ile	Thr	Gln	Thr	275	280	285	
Trp	Gln	Gln	Ala	Asp	Asp	Glu	Asn	Glu	Glu	Ser	Asp	Thr	Pro	Phe	Arg				

RH0020.ST25  
300

<p>290</p> <p>Gln Ile Thr Glu Met Thr Ile Leu Thr Val Gln Leu Ile Val Glu Phe 305 310 315 320</p> <p>Ala Lys Gly Leu Pro Gly Phe Ala Lys Ile Ser Gln Pro Asp Gln Ile 325 330 335</p> <p>Thr Leu Leu Lys Ala Cys Ser Ser Glu Val Met Met Leu Arg Val Ala 340 345 350</p> <p>Arg Arg Tyr Asp Ala Ala Ser Asp Ser Val Leu Phe Ala Asn Asn Gln 355 360 365</p> <p>Ala Tyr Thr Arg Asp Asn Tyr Arg Lys Ala Gly Met Ala Tyr Val Ile 370 375 380</p> <p>Glu Asp Leu Leu His Phe Cys Arg Cys Met Tyr Ser Met Ala Leu Asp 385 390 395 400</p> <p>Asn Ile His Tyr Ala Leu Leu Thr Ala Val Val Ile Phe Ser Asp Arg 405 410 415</p> <p>Pro Gly Leu Glu Gln Pro Gln Leu Val Glu Glu Ile Gln Arg Tyr Tyr 420 425 430</p> <p>Leu Asn Thr Leu Arg Ile Tyr Ile Leu Asn Gln Leu Ser Gly Ser Ala 435 440 445</p> <p>Arg Ser Ser Val Ile Tyr Gly Lys Ile Leu Ser Ile Leu Ser Glu Leu 450 455 460</p> <p>Arg Thr Leu Gly Met Gln Asn Ser Asn Met Cys Ile Ser Leu Lys Leu 465 470 475 480</p> <p>Lys Asn Arg Lys Leu Pro Pro Phe Leu Glu Glu Ile Trp Asp Val Ala 485 490 495</p> <p>Asp Met Ser His Thr Gln Pro Pro Pro Ile Leu Glu Ser Pro Thr Asn 500 505 510</p>	<p>295</p>
---	------------

Leu

<210> 51  
 <211> 4375  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 51  
 tgtaattttg atgggcgccg tgatgcaccg tgtgccatat tgccatccag tcgaatagaa 60  
 aaaaaaaaaa aaaaaaaaaat atcagttgtt ttgtccctcg ctgcctttcg agtgtattcg 120  
 gaatattaga cgtcataatt cacgagtgtc ttttaaattt atatagcgat tagcggggcc 180  
 gtttgttgga cgtgcgcttg cgtttagtgg agtgcaggga tagtgaggcg agtatggtag 240  
 ttcgtggtca tgtcaagtgt ggcgaagaaa gacaagccga cgatgtcggg gacggcgctg 300

RH0020.ST25

atcaactggg	cgcgccgggc	gccgccagge	ccgccgcagc	cgagtcagc	gtcgctgcg	360
ccggcagcca	tgctgcagca	gctcccagc	cagtcaatgc	agtcgttaaa	ccacatccca	420
actgtcgatt	gctcgctcga	tatgcagtgg	cttaatttag	aacctggatt	catgtcgcct	480
atgtcacctc	ctgagatgaa	accagacacc	gccatgcttg	atgggctacg	agacgacgcc	540
acttcgccgc	ctaacttcaa	gaactacccg	cctaataacc	ccctgagtgg	ctccaaacac	600
ctatgctcta	tatggggcga	cagggcgctc	gggaagcact	atgggggtga	cagttgcgaa	660
ggatgcaagg	gtttcttcaa	gcggaccgtc	cggaaggacc	tgctgtacgc	ttgccgggag	720
gagcgggaact	gcatcataga	caagcgacaa	aggaaccgat	gccagtactg	ccgctatcaa	780
aagtgttttg	cttgccgtat	gaagcgagag	gcggtgcaag	aggagcgcca	gaggaatgct	840
cgcgccgcgg	aggatgcgca	cccagtagtc	tcggtgcagg	taagcgatga	gctgtcaatc	900
gagcgcctaa	cggagatgga	gtctttggtg	gcagatccca	gcgaggagtt	ccagttcctc	960
cgcggtggggc	ctgacagcaa	cgtgcctcca	cgttaccgcg	cgcccgcttc	ctccctctgc	1020
caaataaggca	acaagcaaat	agcggcggtt	gtggatatgg	cgcgcgacat	ccctcatttc	1080
gggcagctgg	agctggacga	tcaagtggta	ctcatcaagg	cctcctggaa	tgagctgcta	1140
ctcttcgccca	tcgcctggcg	ctctatggag	tatttggaag	atgagaggga	gaacggggac	1200
ggaacgcgga	gcaccactca	gccacaactg	atgtgtctca	tgcttgcat	gacgttgcac	1260
cgcaactcgg	cgagcaggc	gggcgtgggc	gccatcttcg	accgcgtgct	gtccgagctc	1320
agtctgaaga	tgcgacacct	gcgcattggc	caggccgagt	acgtcgcgct	caaagccatc	1380
gtgctgctca	accctgatgt	gaaaggactg	aagaatcggc	aagaagtga	cgttttgcga	1440
gaaaaaatgt	tctcttgctt	ggagcactac	tgccggcggt	cggaagcaa	cgaggaaggc	1500
cggtttgcgt	ccttgctgct	gcggctgcca	gctctccgct	ccatctcgct	caagagcttc	1560
gaacacctct	acttcttcca	cctcgtggcc	gaaggctcca	tcagcggata	catacgagag	1620
gcgctccgaa	accacgcgcc	tccgatcgac	gtcaatgcca	tgatgtaaag	tgcgatacac	1680
gccctgccga	tgtagaaga	actatggcta	atagaagcga	aactgaatac	atctaggggtg	1740
ggacttaact	tgggactatc	attaaagtat	cacgcaaatt	atgcgtagtc	agaaagtgcg	1800
gtcgatcaaa	cttttttata	aacgaattga	gtttctaacg	actgcaacac	agcggagttt	1860
tgcttctgat	agtttttatt	ctaattggtta	agatgcttta	cacgggcatt	attgacattc	1920
aagtgttaagt	ggaagttagc	aaccttgaca	tttatatcac	gtttgttaatt	ggttaaataa	1980
attaattaat	cacaagtaag	actaacatca	acgtcacgat	actaacgcca	tttagtgata	2040
tttttcatgt	caagaaactc	attgttttga	taaaatat	ttctaattac	tccagtgaac	2100
tcattccaaat	gtgaccagct	ttcccgacga	gttgcccggtg	taaaatcatc	tttagggaca	2160
tatcccccg	tatctcatga	aattccaagg	atcagtaggg	gccaatcccc	ccgatgtgtt	2220
gggaggcaga	attttcgata	atctacgact	attgttagcc	tacgaattag	ttgaattttt	2280



RH0020.ST25

tgaaattatt	tttattaagt	cgccactttc	caaacacatc	agcagggtat	atgtgcaatt	2340
ttgtaacgat	aactctattc	atttctgata	tttatcgaaa	ttttatctta	cataacatgc	2400
tggctggtcc	aggtgtttgg	tagttacata	tgtatctacg	gtttgtttta	aattatagct	2460
tttttattgt	aatctgtata	aaattgagtt	atcttacttc	acactacgat	cgagtaaacc	2520
catcgtcagc	tacgaaaaac	taatcgtata	aggcgtgaaga	gtaaataact	aattgacaac	2580
cagcaacgag	gaccacctca	gtcctcgtgc	ttacattgtg	ccgtagctta	atatgatgga	2640
agctgtcgtc	gttacgacat	tagataaagt	gcatgaatac	caaaaatgta	ccatcccgtg	2700
ctgatctctc	atgctctcgc	tgcgtgggac	ccgtgtcgag	tgtcgtgaagg	actgactaat	2760
attttagact	aggcgtctat	gcttcagtaa	ttccttatac	atattataag	tcattccaaat	2820
aacgagtaag	gcggcatggt	gagatcagca	ttccgagagt	caaagagccc	ctaacgtgac	2880
tgagaagtag	agacaataca	ctgattttct	gagatgaacg	caaccgagat	tgacactaaa	2940
aatctattta	tggatttcaa	aatggcgatg	cttgattgtc	tgcggcgtgg	atagactgaa	3000
atgggtttgc	ttaacactgg	atattgtttt	tattagttaa	tagtcttaca	ttgcaagttg	3060
gtaattcggg	gctaatatcg	accggtttgt	taactatcta	acggttccca	gtgtcaggca	3120
cacatctttc	ccaagcagac	aacgcaagag	tgtacaaaat	gtacatgta	caaaataagg	3180
aacattcgtc	ggataagtgt	aacagttgat	aggtaaagaa	aatggggccg	cctctttatt	3240
attacgtagc	cgtaaaatta	ttaacgtatt	tagtttagat	gttcagctaa	ttaggataat	3300
tctattttgc	gagtacctag	atgtccatag	tgaattaata	taataattag	actgttacgc	3360
gtaggtaatt	ataaagttta	ccaaatctct	cttcaaagca	aaaactttgt	acacttccgt	3420
actgagacgt	cgtagcttat	tctgattcac	gaaatatattg	gatcacattg	ttacaaggcg	3480
accgtcacgt	agtatatgat	tatttacaaa	tgacacgtat	gtatcaatgc	tataagtgtt	3540
ttcgttacat	atgtcgggtg	tttaacgtgc	atttcgatgt	gcagattaaa	aatagcaaga	3600
aatcttgaaa	ttgtttttaga	aaatatattga	tttccttatt	gaaagttatt	tttaaatgta	3660
aatatttcgt	aatcataata	attatgtatt	gtgtagttaa	ttcaccttta	cggttgggat	3720
attatttaat	ggtaggcctac	gaaagtgatt	ataaccatcc	gcgtcctcaa	aaaggccagt	3780
ttatttttgt	acctcataca	tactaattac	gtaagtaata	tcaggcgaat	ggttgactaa	3840
caactaacca	gtattaaaaa	ttaaagact	tcgtcctaata	aaaatgtaata	atctatgtat	3900
aaaaatgaaa	aatctggcgt	ataataggta	aaattaaact	agattgttaa	tgaatgtgat	3960
gtctcataaa	cgtttagttt	ttaatgagaa	acatgttttag	tcgcctacta	taagacgaga	4020
cggcaagctc	accgagttaa	ctcgtaaaca	ggaatgttga	aaaagatgac	acaatttata	4080
tttggatttg	aaattatgac	taacatgcg	ctctatcggt	tgttatggat	gcatagtatt	4140
gctgttgaaa	ataatggaat	taggtaatta	ctgcattaat	gttgaaaact	tgatattatt	4200

RH0020.ST25

ctatggttgg gtatgaattc tatgttgga ggtgtgcagc ggttgtaaag atgatttata 4260  
 atgatgttca ctaaatatct gactaaatgt aagttatattt tttttgtata gacatagctt 4320  
 taagatgaag gtgattaaac tttatcctta tcacaataaa aaaaaaaaaa aaaaaa 4375

<210> 52  
 <211> 472  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 52

Met	Ser	Ser	Val	Ala	Lys	Lys	Asp	Lys	Pro	Thr	Met	Ser	Val	Thr	Ala
1				5					10					15	
Leu	Ile	Asn	Trp	Ala	Arg	Pro	Ala	Pro	Pro	Gly	Pro	Pro	Gln	Pro	Gln
			20					25					30		
Ser	Ala	Ser	Pro	Ala	Pro	Ala	Ala	Met	Leu	Gln	Gln	Leu	Pro	Thr	Gln
		35					40					45			
Ser	Met	Gln	Ser	Leu	Asn	His	Ile	Pro	Thr	Val	Asp	Cys	Ser	Leu	Asp
	50					55					60				
Met	Gln	Trp	Leu	Asn	Leu	Glu	Pro	Gly	Phe	Met	Ser	Pro	Met	Ser	Pro
65				70					75						80
Pro	Glu	Met	Lys	Pro	Asp	Thr	Ala	Met	Leu	Asp	Gly	Leu	Arg	Asp	Asp
				85					90					95	
Ala	Thr	Ser	Pro	Pro	Asn	Phe	Lys	Asn	Tyr	Pro	Pro	Asn	His	Pro	Leu
			100					105					110		
Ser	Gly	Ser	Lys	His	Leu	Cys	Ser	Ile	Cys	Gly	Asp	Arg	Ala	Ser	Gly
		115					120					125			
Lys	His	Tyr	Gly	Val	Tyr	Ser	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe	Lys
	130					135					140				
Arg	Thr	Val	Arg	Lys	Asp	Leu	Ser	Tyr	Ala	Cys	Arg	Glu	Glu	Arg	Asn
145					150					155					160
Cys	Ile	Ile	Asp	Lys	Arg	Gln	Arg	Asn	Arg	Cys	Gln	Tyr	Cys	Arg	Tyr
				165					170					175	
Gln	Lys	Cys	Leu	Ala	Cys	Gly	Met	Lys	Arg	Glu	Ala	Val	Gln	Glu	Glu
			180					185					190		
Arg	Gln	Arg	Asn	Ala	Arg	Gly	Ala	Glu	Asp	Ala	His	Pro	Ser	Ser	Ser
		195					200					205			
Val	Gln	Val	Ser	Asp	Glu	Leu	Ser	Ile	Glu	Arg	Leu	Thr	Glu	Met	Glu
	210					215					220				
Ser	Leu	Val	Ala	Asp	Pro	Ser	Glu	Glu	Phe	Gln	Phe	Leu	Arg	Val	Gly
225					230					235					240
Pro	Asp	Ser	Asn	Val	Pro	Pro	Arg	Tyr	Arg	Ala	Pro	Val	Ser	Ser	Leu

RH0020.ST25

245 250 255  
 Cys Gln Ile Gly Asn Lys Gln Ile Ala Ala Leu Val Val Trp Ala Arg  
 260 265 270  
 Asp Ile Pro His Phe Gly Gln Leu Glu Leu Asp Asp Gln Val Val Leu  
 275 280 285  
 Ile Lys Ala Ser Trp Asn Glu Leu Leu Leu Phe Ala Ile Ala Trp Arg  
 290 295 300  
 Ser Met Glu Tyr Leu Glu Asp Glu Arg Glu Asn Gly Asp Gly Thr Arg  
 305 310 315 320  
 Ser Thr Thr Gln Pro Gln Leu Met Cys Leu Met Pro Gly Met Thr Leu  
 325 330 335  
 His Arg Asn Ser Ala Gln Gln Ala Gly Val Gly Ala Ile Phe Asp Arg  
 340 345 350  
 Val Leu Ser Glu Leu Ser Leu Lys Met Arg Thr Leu Arg Met Asp Gln  
 355 360 365  
 Ala Glu Tyr Val Ala Leu Lys Ala Ile Val Leu Leu Asn Pro Asp Val  
 370 375 380  
 Lys Gly Leu Lys Asn Arg Gln Glu Val Asp Val Leu Arg Glu Lys Met  
 385 390 395 400  
 Phe Ser Cys Leu Asp Asp Tyr Cys Arg Arg Ser Arg Ser Asn Glu Glu  
 405 410 415  
 Gly Arg Phe Ala Ser Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser Ile  
 420 425 430  
 Ser Leu Lys Ser Phe Glu His Leu Tyr Phe Phe His Leu Val Ala Glu  
 435 440 445  
 Gly Ser Ile Ser Gly Tyr Ile Arg Glu Ala Leu Arg Asn His Ala Pro  
 450 455 460  
 Pro Ile Asp Val Asn Ala Met Met  
 465 470

<210> 53  
 <211> 1404  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> misc\_feature  
 <223> Novel Sequence

<400> 53  
 atggacacca aacatttcct gccgctcgac ttctctaccc aggtgaactc ttcgccctc 60  
 aactctccaa cgggtcgagg ctccatggct gtcccctcgc tgcaccctc cttgggtccg 120  
 ggaatcggct ctccactggg ctgcctggg cagctgcact ctctatcag caccctgagc 180  
 tcccccatca atggcatggg tccgcccttc tctgtcatca gctcccccat ggcccgcac 240  
 tccatgtcgg taccacaccac accacattg ggcttcggga ctggtagccc ccagctcaat 300

RH0020.ST25

```

tcacccatga accctgtgag cagcactgag gatatacaagc cgccactagg cctcaatggc 360
gtcctcaagg ttccctgccca tccctcagga aatatggcct ccttcaccaa gcacatctgt 420
gctatctgtg gggaccgctc ctccaggcaaa cactatgggg tatacagttg tgagggctgc 480
aagggcttct tcaagaggac agtacgcaaa gacctgacct acacctgccg agacaacaag 540
gactgcctga tcgacaagag acagcggaac cgggtgtcagt actgccgcta ccagaagtgc 600
ctggccatgg gcatgaagcg ggaagctgtg caggaggagc ggcagcgggg caaggaccgg 660
aatgagaacg aggtggagtc caccagcagt gccaacgagg acatgcctgt agagaagatt 720
ctggaagccg agcttgctgt cgagcccaag actgagacat acgtggaggc aaacatgggg 780
ctgaacccca gctcaccaaa tgaccctggt accaacaatct gtcaagcagc agacaagcag 840
ctcttcaact ttgtggagtg ggccaagagg atcccacact tttctgagct gcccctagac 900
gaccaggtca tcctgctacg ggcaggtctg aacgagctgc tgatcgctc cttctccac 960
cgctccatag ctgtgaaaga tgggattctc ctggccaccg gcctgcacgt acaccggaac 1020
agcgctcaca gtgctggggg gggcgccatc tttgacaggg tgctaacaga gctgggtgtct 1080
aagatgcgtg acatgcagat ggacaagaag gagctgggct gcctgcgagc cattgtcctg 1140
ttcaaccctg actctaaggg gctctcaaac cctgctgagg tggaggcgtt gagggagaag 1200
gtgtatgcgt cactagaagc gtactgcaaa cacaagtacc ctgagcagcc gggcaggttt 1260
gccaaactgc tgctccgctt gcctgcaact cgttccatcg ggctcaagtg cctggagcac 1320
ctgttcttct tcaagctcat cggggacacg cccatcgaca ccttcctcat ggagatgctg 1380
gaggcaccac atcaagccac ctg 1404

```

<210> 54  
 <211> 467  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <221> misc feature  
 <223> Novel Sequence

<400> 54

```

Met Asp Thr Lys His Phe Leu Pro Leu Asp Phe Ser Thr Gln Val Asn
1          5          10          15
Ser Ser Ser Leu Asn Ser Pro Thr Gly Arg Gly Ser Met Ala Val Pro
20          25          30
Ser Leu His Pro Ser Leu Gly Pro Gly Ile Gly Ser Pro Leu Gly Ser
35          40          45
Pro Gly Gln Leu His Ser Pro Ile Ser Thr Leu Ser Ser Pro Ile Asn
50          55          60
Gly Met Gly Pro Pro Phe Ser Val Ile Ser Ser Pro Met Gly Pro His
65          70          75          80

```

RH0020.ST25

Ser Met Ser Val Pro Thr Thr Pro Thr Leu Gly Phe Gly Thr Gly Ser  
 85 90 95  
 Pro Gln Leu Asn Ser Pro Met Asn Pro Val Ser Ser Thr Glu Asp Ile  
 100 105 110  
 Lys Pro Pro Leu Gly Leu Asn Gly Val Leu Lys Val Pro Ala His Pro  
 115 120 125  
 Ser Gly Asn Met Ala Ser Phe Thr Lys His Ile Cys Ala Ile Cys Gly  
 130 135 140  
 Asp Arg Ser Ser Gly Lys His Tyr Gly Val Tyr Ser Cys Glu Gly Cys  
 145 150 155 160  
 Lys Gly Phe Phe Lys Arg Thr Val Arg Lys Asp Leu Thr Tyr Thr Cys  
 165 170 175  
 Arg Asp Asn Lys Asp Cys Leu Ile Asp Lys Arg Gln Arg Asn Arg Cys  
 180 185 190  
 Gln Tyr Cys Arg Tyr Gln Lys Cys Leu Ala Met Gly Met Lys Arg Glu  
 195 200 205  
 Ala Val Gln Glu Glu Arg Gln Arg Gly Lys Asp Arg Asn Glu Asn Glu  
 210 215 220  
 Val Glu Ser Thr Ser Ser Ala Asn Glu Asp Met Pro Val Glu Lys Ile  
 225 230 235 240  
 Leu Glu Ala Glu Leu Ala Val Glu Pro Lys Thr Glu Thr Tyr Val Glu  
 245 250 255  
 Ala Asn Met Gly Leu Asn Pro Ser Ser Pro Asn Asp Pro Val Thr Asn  
 260 265 270  
 Ile Cys Gln Ala Ala Asp Lys Gln Leu Phe Thr Leu Val Glu Trp Ala  
 275 280 285  
 Lys Arg Ile Pro His Phe Ser Glu Leu Pro Leu Asp Asp Gln Val Ile  
 290 295 300  
 Leu Leu Arg Ala Gly Trp Asn Glu Leu Leu Ile Ala Ser Phe Ser His  
 305 310 315 320  
 Arg Ser Ile Ala Val Lys Asp Gly Ile Leu Leu Ala Thr Gly Leu His  
 325 330 335  
 Val His Arg Asn Ser Ala His Ser Ala Gly Val Gly Ala Ile Phe Asp  
 340 345 350  
 Arg Val Leu Thr Glu Leu Val Ser Lys Met Arg Asp Met Gln Met Asp  
 355 360 365  
 Lys Thr Glu Leu Gly Cys Leu Arg Ala Ile Val Leu Phe Asn Pro Asp  
 370 375 380  
 Ser Lys Gly Leu Ser Asn Pro Ala Glu Val Glu Ala Leu Arg Glu Lys  
 385 390 395 400  
 Val Tyr Ala Ser Leu Glu Ala Tyr Cys Lys His Lys Tyr Pro Glu Gln  
 405 410 415  
 Pro Gly Arg Phe Ala Lys Leu Leu Leu Arg Leu Pro Ala Leu Arg Ser

RH0020.ST25

420 425 430  
 Ile Gly Leu Lys Cys Leu Glu His Leu Phe Phe Phe Lys Leu Ile Gly  
 435 440 445  
 Asp Thr Pro Ile Asp Thr Phe Leu Met Glu Met Leu Glu Ala Pro His  
 450 455 460  
 Gln Ala Thr  
 465  
 <210> 55  
 <211> 309  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <221> misc feature  
 <223> Novel Sequence  
 <400> 55  
 ggtgtggaaa gtccccaggc tccccagcag gcagaagtat gcaaagcatg catctcaatt 60  
 agtcagcaac caggtgtgga aagtccccag gctccccagc aggcagaagt atgcaaagca 120  
 tgcattctcaa ttagtcagca accatagtcc cgcccctaac tccgcccata ccgcccctaa 180  
 ctccgcccag ttccgcccata tctccgcccc atggctgact aaaaaaaaaa atttatgcag 240  
 aggccgaggg cgccctcgccc tctgagctat tccagaagta gtgaggaggc ttttttggag 300  
 gcctaggct 309  
 <210> 56  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <221> misc feature  
 <223> Novel Sequence  
 <400> 56  
 tatataatgg atccccgggt accg 24  
 <210> 57  
 <211> 1653  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <221> misc feature  
 <223> Novel Sequence  
 <400> 57  
 atggaagacg caaaaaacat aaagaaaggc cgggcgccat tctatcctct agaggatgga 60  
 accgctggag agcaactgca taaggctatg aagagatacg ccctgggttc tggaacaatt 120  
 gcttttacag atgcacatat cgaggtgaac atcacgtacg cggaataact cgaaatgtcc 180

RH0020.ST25

```

gttcggttgg cagaagctat gaaacgatat gggctgaata caaatcacag aatcgtcgta 240
tgcagtga aa actctcttca attctttatg ccggtgttgg gcgcgttatt tatcggagtt 300
gcagttgcgc ccgcgaacga catttataat gaacgtgaat tgctcaacag tatgaacatt 360
tcgcagccta ccgtagtggt tgtttccaaa aagggttgc aaaaaatttt gaacgtgcaa 420
aaaaaattac caataatcca gaaaattatt atcatggatt ctaaaacgga ttaccaggga 480
tttcagtcga tgtacacggt cgtcacatct catctacctc ccggttttaa tgaatacga 540
tttgtaccag agtcctttga tcgtgacaaa acaattgcac tgataatgaa ttctcttgga 600
tctactgggt tacctaagggt tgtggccctt ccgcatagaa ctgcctgcgt cagattctcg 660
catgccagag atcctatttt tggcaatcaa atcattccgg atactgcgat ttttaagtgtt 720
gttccattcc atcacggttt tggaatgttt actacactcg gatatttgat atgtggattt 780
cgagtcgtct taatgtatag atttgaagaa gagctgtttt tacgatccct tcaggattac 840
aaaattcaaa gtgcgttgct agtaccaacc ctattttcat tcttcgcaa aagcactctg 900
attgacaaat acgatttatc taatttacac gaaattgctt ctgggggcgc acctctttcg 960
aaagaagtcg gggaagcgggt tgcaaacgc ttccatcttc cagggatacg acaaggatat 1020
gggctcactg agactacatc agctattctg attacacccg agggggatga taaaccgggc 1080
gcggtcggta aagttgttcc attttttgaa gcgaagggtg tggatctgga taccgggaaa 1140
acgctgggcg ttaatcagag aggcgaatta tgtgtcagag gacctatgat tatgtccggt 1200
tatgtaaaca atccggaagc gaccaacgcc ttgattgaca aggatggatg gctacattct 1260
ggagacatag ctactggga cgaagacgaa cacttcttca tagttgaccg cttgaagtct 1320
ttaattaaat acaaaggata tcaggtggcc ccgctgaat tggaatcgat attgttacia 1380
caccccaaca tcttcgacgc gggcgtggca ggtcttcccg acgatgacgc cgggtgaactt 1440
cccgccgccg ttgttgtttt ggagcacgga aagacgatga cggaaaaaga gatcgtggat 1500
tacgtcgcca gtcaagtaac aaccgcgaaa aagttgcgcg gaggagtgtt gtttgtggac 1560
gaagtaccga aaggtcttac cggaaaactc gacgcaagaa aaatcagaga gatcctcata 1620
aaggccaaga agggcgga aa gtccaaattg taa 1653

```

```

<210> 58
<211> 867
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc_feature
<223> Novel Sequence

```

```

<400> 58
aagcgagagg cgggtgcaaga ggagcgccag aggaatgctc gcggcgcgga ggatgcgcac 60

```

## RH0020.ST25

```

ccgagtagct cggtgcaggt aagcgatgag ctgtcaatcg agcgcctaac ggagatggag 120
tcttttggtgg cagatcccag cgaggagttc cagttcctcc gcgtaggggcc tgacagcaac 180
gtgcctccac gtiaccgcgc gcccgctctc tccctctgcc aaataggcaa caagcaaata 240
gcggcggttgg tggatatgggc gcgcgacatc cctcatttcg ggcagctgga gctggacgat 300
caagtggtag tcatcaaggc ctcttggaat gagctgtac tcttcgccat cgcctggcgc 360
tctatggagt atttggaaga tgagagggag aacggggacg gaacgcggag caccactcag 420
ccacaactga tgtgtctcat gcctggcatg acgttgaccc gcaactcggc gcagcaggcg 480
ggcgtggggcg ccatcttcga ccgcgtgctg tccgagctca gtctgaagat gcgcaccttg 540
cgcatggacc aggcggagta cgtcgcgctc aaagccatcg tgctgctcaa ccctgatgtg 600
aaaggactga agaatcggca agaagttgac gttttgcgag aaaaaatgtt ctcttgccctg 660
gacgactact gccggcggtc gcgaagcaac gaggaaggcc ggtttgctgc cttgctgctg 720
cggctgccag ctctccgctc catctcgtc aagagcttcg aacacctcta cttcttccac 780
ctcgtggccg aaggctccat cagcggatac atacgagagg cgctccgaaa ccacgcgcct 840
ccgatcgacg tcaatgccat gatgtaa 867

```

```

<210> 59
<211> 225
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc feature
<223> Novel Sequence

```

```

<400> 59
tcgacattgg acaagtgcac tgaacccttg tctctcgaga gacaaggggg ttcaatgcac 60
ttgtccaatg tcgagagaca aggggggttca atgcacttgt ccaatgtcga gagacaaggg 120
ggttcaatgc acttgtccaa tgtcgagaga caaggggggt caatgcactt gtccaatgtc 180
gagagacaag ggggttcaat gcacttgtcc aatgtcgact ctaga 225

```

```

<210> 60
<211> 619
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc feature
<223> Novel Sequence

```

```

<400> 60
cgttacataa cttacggtaa atggcccgcg tggctgaccg cccaacgacc cccgcccatt 60
gacgtcaata atgacgtatg ttcccatagt aacgccaaata gggactttcc attgacgtca 120
atgggtggag tattttacggt aaactgcccc cttggcagta catcaagtgt atcatatgcc 180

```



## RH0020.ST25

```

aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt atgcccagta 240
catgacctta tgggactttc ctacttggca gtacatctac gtattagtca tcgctattac 300
catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg actcacgggg 360
atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc aaaatcaacg 420
ggactttcca aaatgtcgta acaactccgc cccattgacg caaatgggcg gtaggcgtgt 480
acgggtgggag gtctatataa gcagagctcg tttagtgaac cgtcagatcg cctggagacg 540
ccatccacgc tgttttgacc tccatagaag acaccgggac cgatccagcc tccgcgcccg 600
ggaacggtgc attggaacg 619

```

```

<210> 61
<211> 262
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc_feature
<223> Novel Sequence

```

```

<400> 61
atgtagtctt atgcaatact cttgtagtct tgcaacatgg taacgatgag ttagcaacat 60
gccttacaag gagagaaaaa gcaccgtgca tgccgatagg tggaagtaag gtggtacgat 120
cgtgccttat taggaaggca acagacgggt ctgacatgga ttggacgaac cactgaattc 180
cgcattgcag agatattgta ttttaagtgc tagctcgata caataaacgc catttgacca 240
ttcaccacat tggagtgcac ct 262

```

```

<210> 62
<211> 1247
<212> DNA
<213> Artificial Sequence

```

```

<220>
<221> misc_feature
<223> Novel Sequence

```

```

<400> 62
tctatttcct caggccgtga ggaactgtcg ccagcttcaa gtataaatgg gtgcagtaca 60
gatggcgagg caccagctca gaagaagggc cctgcgcccc gtcagcaaga ggaactgtgt 120
ctggtatgcg gggacagagc ctccggatac cactacaatg cgctcacgtg tgaaggggtgt 180
aaagggttct tcagacggag tgttaccaaa aatgcggttt atatttgtaa attcggtcac 240
gcttgcgaaa tggacatgta catgcgacgg aaatgccagg agtgccgcct gaagaagtgc 300
ttagctgtag gcatgaggcc tgagtgcgta gtacccgaga ctcagtgcgc catgaagcgg 360
aaagagaaga aagcacagaa ggagaaggac aaactgcctg tcagcacgac gacggtggac 420

```

RH0020.ST25

```

gaccacatgc cgcccattat gcagtgtaa cctccacctc ctgaagcagc aaggattcac 480
gaagtgggtcc caaggtttct ctccgacaag ctgttgagaga caaaccggca gaaaaacatc 540
ccccagttga cagccaacca gcagttcctt atcgccaggc tcattctggta ccaggacggg 600
tacgagcagc cttctgatga agatttgaag aggattacgc agacgtggca gcaagcggac 660
gatgaaaacg aagagtctga cactcccttc cgccagatca cagagatgac tatcctcacg 720
gtccaactta tcgtggagtt cgcgaaagga ttgccagggt tcgccaagat ctgcgagcct 780
gatcaaatta cgctgcttaa ggcttgctca agtgaggtaa tgatgctccg agtcgcgcga 840
cgatacgatg cggcctcaga cagtgttctg ttgcggaaca accaagcgta cactcgcgac 900
aactaccgca aggctggcat ggcctacgtc atcgaggatc tactgcactt ctgccgggtgc 960
atgtactcta tggcgttgga caacatccat tacgcgctgc tcacggctgt cgtcatcttt 1020
tctgaccggc cagggttgga gcagccgcaa ctggtggaag aaatccagcg gtactacctg 1080
aatagctcc gcatttatat cctgaaccag ctgagcgggt cggcgcgctt gtccgtcata 1140
tacggcaaga tcctctcaat cctctctgag ctacgcacgc tcggcatgca aaactccaac 1200
atgtgcatct cctcaagct caagaacaga aagctgccgc ctttct 1247

```

```

<210> 63
<211> 440
<212> PRT
<213> Artificial Sequence

```

```

<220>
<221> misc feature
<223> Novel Sequence

```

```

<400> 63

```

```

Ser Ile Ser Ser Gly Arg Glu Glu Leu Ser Pro Ala Ser Ser Ile Asn
1          5          10          15
Gly Cys Ser Thr Asp Gly Glu Ala Arg Arg Gln Lys Lys Gly Pro Ala
20          25          30
Pro Arg Gln Gln Glu Glu Leu Cys Leu Val Cys Gly Asp Arg Ala Ser
35          40          45
Gly Tyr His Tyr Asn Ala Leu Thr Cys Glu Gly Cys Lys Gly Phe Phe
50          55          60
Arg Arg Ser Val Thr Lys Asn Ala Val Tyr Ile Cys Lys Phe Gly His
65          70          75          80
Ala Cys Glu Met Asp Met Tyr Met Arg Arg Lys Cys Gln Glu Cys Arg
85          90          95
Leu Lys Lys Cys Leu Ala Val Gly Met Arg Pro Glu Cys Val Val Pro
100         105         110
Glu Thr Gln Cys Ala Met Lys Arg Lys Glu Lys Lys Ala Gln Lys Glu
115         120         125

```

RH0020.ST25

Lys Asp Lys Leu Pro Val Ser Thr Thr Thr Val Asp Asp His Met Pro  
 130 135 140  
 Pro Ile Met Gln Cys Glu Pro Pro Pro Pro Glu Ala Ala Arg Ile His  
 145 150 155 160  
 Glu Val Val Pro Arg Phe Leu Ser Asp Lys Leu Leu Glu Thr Asn Arg  
 165 170 175  
 Gln Lys Asn Ile Pro Gln Leu Thr Ala Asn Gln Gln Phe Leu Ile Ala  
 180 185 190  
 Arg Leu Ile Trp Tyr Gln Asp Gly Tyr Glu Gln Pro Ser Asp Glu Asp  
 195 200 205  
 Leu Lys Arg Ile Thr Gln Thr Trp Gln Gln Ala Asp Asp Glu Asn Glu  
 210 215 220  
 Glu Ser Asp Thr Pro Phe Arg Gln Ile Thr Glu Met Thr Ile Leu Thr  
 225 230 235 240  
 Val Gln Leu Ile Val Glu Phe Ala Lys Gly Leu Pro Gly Phe Ala Lys  
 245 250 255  
 Ile Ser Gln Pro Asp Gln Ile Thr Leu Lys Ala Cys Ser Ser Glu  
 260 265 270  
 Val Met Met Leu Arg Val Ala Arg Arg Tyr Asp Ala Ala Ser Asp Ser  
 275 280 285  
 Val Leu Phe Ala Asn Asn Gln Ala Tyr Thr Arg Asp Asn Tyr Arg Lys  
 290 295 300  
 Ala Gly Met Ala Tyr Val Ile Glu Asp Leu Leu His Phe Cys Arg Cys  
 305 310 315 320  
 Met Tyr Ser Met Ala Leu Asp Asn Ile His Tyr Ala Leu Leu Thr Ala  
 325 330 335  
 Val Val Ile Phe Ser Asp Arg Pro Gly Leu Glu Gln Pro Gln Leu Val  
 340 345 350  
 Glu Glu Ile Gln Arg Tyr Tyr Leu Asn Thr Leu Arg Ile Tyr Ile Leu  
 355 360 365  
 Asn Gln Leu Ser Gly Ser Ala Arg Ser Ser Val Ile Tyr Gly Lys Ile  
 370 375 380  
 Leu Ser Ile Leu Ser Glu Leu Arg Thr Leu Gly Met Gln Asn Ser Asn  
 385 390 395 400  
 Met Cys Ile Ser Leu Lys Leu Lys Asn Arg Lys Leu Pro Pro Phe Leu  
 405 410 415  
 Glu Glu Ile Trp Asp Val Ala Asp Met Ser His Thr Gln Pro Pro Pro  
 420 425 430  
 Ile Leu Glu Ser Pro Thr Asn Leu  
 435 440

&lt;210&gt; 64

&lt;211&gt; 943

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

RH0020.ST25

&lt;220&gt;

&lt;221&gt; misc feature

&lt;223&gt; Novel Sequence

&lt;400&gt; 64

atgacttcga aagtttatga tccagaacaa aggaaacgga tgataactgg tccgcagtgg	60
tgggccagat gtaaacaat gaatgttctt gattcattta ttaattatta tgattcagaa	120
aaacatgcag aaaatgctgt tatTTTTTTta catggtaacg cggcctcttc ttatttatgg	180
cgacatgttg tgccacatat tgagccagta gcgcggtgta ttataccaga ccttattggt	240
atggggcaaat caggcaaact tggtaatggt tcttataggt tacttgatca ttacaaatat	300
cttactgcat ggtttgaact tcttaattta ccaaagaaga tcatttttgt cggccatgat	360
tggggtgctt gtttggcatt tcattatagc tatgagcatc aagataagat caaagcaata	420
gttcacgctg aaagtgtagt agatgtgatt gaatcatggg atgaatggcc tgatattgaa	480
gaagatatatg cgttgatcaa atctgaagaa ggagaaaaaa tggtttttga gaataacttc	540
ttcgtggaaa ccatgttgcc atcaaaaatc atgagaaagt tagaaccaga agaatttgca	600
gcatatcttg aaccattcaa agagaaaggt gaagttcgtc gtccaacatt atcatggcct	660
cgtgaaatcc cgttagttaa aggtggtaaa cctgacgttg tacaaattgt taggaattat	720
aatgcttatc tacgtgcaag tgatgattta ccaaaaatgt ttattgaatc ggaccagga	780
ttcttttcca atgctattgt tgaagggtgcc aagaagtttc ctaatactga atttgtcaaa	840
gtaaaaggtc ttcatTTTTc gcaagaagat gcacctgatg aaatgggaaa atatatcaaa	900
tcgttcgttg agcgagttct caaaaatgaa caataattct aga	943